

CHRISTIAN CAPAPÉ¹, ABDERRAHMANE KASSAR²,
CHRISTIAN REYNAUD³, FARID HEMIDA²

¹ Laboratoire d'Ichtyologie, Université de Montpellier, case 104,
34095 Montpellier cedex 5, France

² École Nationale Supérieure des Sciences de la Mer et de l'Aménagement du Littoral
(ENSSMAL), BP 19, Bois des Cars, 16320 Dely Ibrahim, Algiers, Algeria

³ LIRDEF, E. A. 3749, Faculté d'Éducation, Université Montpellier, 2, place Marcel Godechot,
B.P. 4152, 34092 Montpellier cedex5, France
email: capape@univ-montp2.fr

ATYPICAL CHARACTERISTICS IN BLACKMOUTH CATSHARK, *GALEUS MELASTOMUS* (CHONDRICHTHYES: SCYLIORHINIDAE) FROM THE ALGERIAN COAST (SOUTHERN MEDITERRANEAN SEA)

SUMMARY

The authors report in the present paper the capture of an abnormal specimen of the blackmouth catshark *Galeus melastomus* RAFINESQUE, 1810 from the central coast of Algeria (southern Mediterranean Sea). The specimen was a sub-mature female having 478 mm in total length (TL) and weighing 265 g in total body weight (TBW). The specimen displayed a severe atrophy of the right clasper and a poor development of the right pelvic fin. These abnormalities are compared with other similar patterns reported in elasmobranch. The origin of the abnormalities remains questionable, due to unfavourable environmental conditions (action of pollutants) or to endogenous origin or genetic during embryonic development.

INTRODUCTION

Blackmouth catshark, *Galeus melastomus* Rafinesque, 1810 is a small shark species known from the north-eastern Atlantic, the Bay of Biscay to the coast of Portugal (QUÉRO *et al.*, 2003). South the Strait of Gibraltar, it occurs off Morocco (LORIS and RUCABADO, 1998), but it is replaced by its close relative species *G. polli* CADENAT, 1959 from the coast of Mauritania (MAURIN and BONNET, 1970), to the Gulf of Guinea (BLACHE *et al.*, 1970).

Galeus melastomus is known in the entire Mediterranean Sea (CAPAPÉ, 1989; SERENA, 2005), including the Adriatic Sea (LIPEJ and DULCIC, 2010) and the eastern Levant Basin (GOLANI, 2005). The species occurs off the Maghreb shore, where it is rather captured at depths over 100-350 m approximatively, MOROCCO (LLORIS and RUCABADO, 1998), ALGERIA (HEMIDA, 2005; AÏT DARNA *et al.*, 2018), TUNISIA (BRADAI *et al.*, 2004). Following CAPAPÉ *et al.* (2008), *G. melastomus* formed the object of various studies concerning some traits of its distribution, size-structure, reproductive biology, and food and feeding habits from the Mediterranean and outside this sea.

Investigations regularly conducted in the central region of the Algerian coast offered us the opportunity to collect several specimens of *Galeus melastomus*, and among these specimens we have found a male displaying an abnormality of pelvic fins and claspers. This abnormality is described and commented in the present paper.

MATERIALS AND METHODS

A total of 211 specimens of *Galeus melastomus* were collected during trawl surveys from the central region of the Algerian coast. Of the 102 males, a single specimen exhibiting an abnormality was captured off Ain Biane (Fig. 1), on 14 July 2015, by $36^{\circ} 48' 39''$ and $2^{\circ} 51' 32''$ E, at a depth of 350 m approximatively, on sandy-muddy bottom, together with teleost species such as bluemouth rockfish *Helicolenus dactylopterus* (Delaroche, 1809), black anglerfish *Lophius budegassa* (Spinola, 1807) and greater fork-beared *Phycis blennoides* (Brünnich, 1768).




Fig. 1. Map of the Algerian coast indicating the capture site of the abnormal specimen of *Galeus melastomus* (black star).

The collected specimens were measured for total length (TL) recorded to the nearest millimetre and weighed for total body weight (TBW), recorded to the nearest gram. The 102 sampled males ranged between 252 and 562 mm TL and weighed between 44 and 480 g (TBW). All fresh specimens were

measured to nearest millimetre and weighed to nearest gram. However, a sub-sample of 58 normal specimens was used in order to obtain a more visible view, allowing to clearly distinguish the abnormal specimen from the normal specimens.

The relation between the total length (TL) and the total body weight (TBW) was used as a complement following FROESE *et al.* (2011), all specimens, normal and abnormal to show if this latter is able to develop in the wild as normal specimens. This LWR is $TBW = aTL^b$, and was converted into its linear regression, expressed in decimal logarithmic coordinates and correlations were assessed by least-squares regression. as: $\log TBW = \log a + b \log TL$: Significance of constant b differences was assessed to the hypothesis of isometric growth if $b = 3$, positive allometry if $b > 3$, negative isometry if $b < 3$ (PAULY, 1983). Comparison of means were carried out by using *t*-test. These two latter tests were performed by using logistic model STATVIEW 5.0.

RESULTS AND DISCUSSION

All sampled specimens were identified as *Galeus melastomus* via the combination of main morphological characters, as follows: snout longer than width across the mouth, nostrils widely separated from mouth, first dorsal fin originating over rear part of pelvic fins, a crest of margin of modified denticles on anterior part of caudal, dark-brown blotches along back and sides. These characters are in total accordance with SPRINGER (1979), QUÉRO (1984) and QUÉRO *et al.* (2003).

The abnormal specimen was a male which measured 478 mm TL and weighed 265 g TBW. The left pelvic fin was completely developed and slightly larger than the left clasper which was not elongated, rigid, probably not calcified (Fig. 2). The morphology of this clasper is characteristic of a sub-adult stage described by CAPAPÉ *et al.* (2008) in *G. melastomus*. The right pelvic fin is obviously smaller than the left pelvic fin and exhibited on its internal margin rather appeared as a roll-shaped structure rather than an inconspicuous clasper (Fig. 3). Therefore, we assume that the clasper is severely atrophied in this specimen male of *G. melastomus*.

Size at first maturity and maximum size reported in *G. melastomus* from some Mediterranean areas are summarized in Table 1. It suggests that the present specimen was probably a sub-adult specimen and such statement is corroborated by the morphology of the left clasper.

Abnormalities of claspers are reported in some elasmobranch species as lack or atrophy of this copulatory organ, and are included in Table 2. It clearly appears all recorded instances concerned batoid species which generally live close or buried in soft bottoms. These habitats faced to a large expo-




Fig. 2. Abnormal specimen of *Galeus melastomus*. A. Dorsal surface. B. Ventral surface. Scale bar = 100 mm.




Fig. 3. Abnormal specimen of *Galeus melastomus*. L Cl: left clasper. L Pel Ffin: left pelvic fin. RCL: right clasper. R Pel Fin: right pelvic fin. Scale bar = 50 mm.

Area	Size at sexual maturity (TL, mm)		Maximum size (TL, mm)		Authors
	Males	Females	Males	Females	
Tunisian coast	> 420	390-420	560	550	CAPAPÈ and ZAOUALI (1977)
Ionian Sea	450	490	510	550	TURSI <i>et al.</i> (1993)
Adriatic Sea	450	490	-	580	UNGARO <i>et al.</i> (1996)
Tyrrhenian Sea	-	-	500	500	RINELLI <i>et al.</i> (2005)
Languedoc Coast	510-550	520-610	620	640	CAPAPÈ <i>et al.</i> (2008)
Eastern Algeria	540	540	560	560	AIT DARNA <i>et al.</i> (2018)

Table 1. Size at sexual maturity and maximal size of *Galeus melastomus* recorded from some Mediterranean areas.

Species	Region	Condition of claspers	Authors
<i>Raja miraletus</i> (Linnaeus, 1758)	Tunisian coast	Lack of left clasper	QUIGNARD and CAPAPÈ (1972)
<i>Bathyraja interrupta</i> (Gill and Townsend, 1897)	Coast of Alaska	Atrophy of both claspers	HAAS and EBERT (2008)
<i>Pteroplatytrygon violacea</i> (Bonaparte, 1832)	Brazilian coast	Lack of left clasper	RIBEIRO-PRADO <i>et al.</i> (2009)
<i>Dasyatis tortonesei</i> Capapè, 1975	Tunisian coast	Atrophy of both claspers	CAPAPÈ <i>et al.</i> (2012)
<i>Urotrygon microphthalmum</i> Delsman, 1941	Brazilian coast	Atrophy of left clasper, lack of right clasper	SANTANDER-NETO and LESSA (2013)
<i>Urotrygon chilensis</i> (Günther, 1872)	West coast of Mexico	Lack of left clasper	TORRES-HUERTA <i>et al.</i> (2015)
<i>Zapteryx exasperata</i> (Jordan and Gilbert, 1880)	West coast of Mexico	Lack of right clasper, atrophy of left pelvic fin	GONZÁLEZ-GONZÁLEZ <i>et al.</i> (2016)
<i>Myliobatis aquila</i> (Linnaeus, 1758)	Tunisian coast	Atrophy of both claspers	RAFRAFI- <i>et al.</i> (2017)
<i>Pseudobathos percellens</i> (Walbaum, 1792)	Caribbean Sea	Lack of left clasper, atrophy of right pelvic fin	EHEMANN and GONZÁLEZ- GONZÁLEZ (2018)
<i>Galeus melastomus</i> Rafinesque, 1810	Algerian coast	Severe atrophy of right clasper and pelvic fin	This study

Table 2. Condition of clasper and pelvic fin (lack and/or atrophy) observed in *Galeus melastomus* collected in some marine regions.

sure to radioactive contamination (YANO and TANAKA, 1989) and to pollutant dumps with high values of heavy metals concentrations (RIBEIRA-PRADO *et al.*, 2008) and chemical particulates such as DDT (TORRES-HUERTA *et al.*, 2015). *G. melastomus* is a benthic species but also an oviparous species which lays its egg cases directly in the wild, and embryos are not protected by mother's uteri as for viviparous species and underget the unfavourable environmental parameters (RIBEIRA-PRADO *et al.*, 2008). The study area where our sample was collected is located in the central region of Algeria which faced to different categories of pollutants such as heavy metals (BENAMAR *et al.*, 1999) and polychlorinated biphenyls (FOUIAL-DJEBBAR *et al.*, 2011)

On the other hand, the specimens displaying some abnormalities and reported in the present paper live in tropical and equatorial seas. The waters of the Algerian coast are becoming warmer as for the entire Mediterranean Sea (FRANCOUR *et al.*, 1994), and as a consequence they are progressively invaded by non-indigenous species (HEMIDA *et al.*, 2018). However, the role of temperature in the development of such abnormalities remains only a possible hypothesis which needs adequate studies, but it cannot be totally ruled out. Among the causes of these reproductive abnormalities, lack or atrophy of claspers, EHEMANN and GONZÁLEZ-GONZÁLEZ (2018) noted the most probable origin is related to the embryonic development. This opinion is in total accordance with BENSAM (1965) who emphasized that such deformities are probably caused by intrauterine pressure exerted by other embryos in viviparous species. Conversely, BONFIL (1989) suggested that the pre-natal abnormalities have a genetic origin or related to mutations.

The lack of one clasper probably plays a minor role in reproduction process due to the fact that male specimens introduce a single clasper during copulation (CHAPMAN *et al.*, 2003). Conversely, atrophy of both claspers reduces the success of the reproduction process. The relationship TBW vs TL including the abnormal specimen and other specimens of similar sizes from our sample of *G. melastomus* is $\log TWB = -5.429 + 2.94 * \log TL$; $r = 0.97$; $n = 58$, displaying an allometry slightly negative (Fig. 4), all specimens having a regular increase in development. Therefore, both atrophies of right clasper and right pelvic fin did not assume the development of the abnormal specimen in the wild, similar patterns were provided in two other batoid species (CAPAPÉ *et al.*, 2015a, b).




Fig. 4. Relationship total body mass (TBM) versus disc width (TL) expressed in logarithmic coordinates for normal and abnormal specimens of *Galeus melastomus* collected from the Algerian waters.

REFERENCES

- AÏT DARNA S., TALAB BENDIAB A.A., MOUFFLOK S., BELMAHI A.-E., BOUDERBALA M., 2018. Observations on distribution, biology, growth, diet and feeding strategy of black-mouth catshark, *Galeus melastomus* (RAFINESQUE, 1810) (Chondrichthyes Scyliorhinidae) in western Algerian coast. *Biodiversity Records*, **9** (4): 357-366. DOI: 10.31396/Biodiv.Jour.2018.9.4.357.368.
- BENAMAR M.A., TOUMERT I., TOBBECHE S., TCHANTCHANE A., CHALABI A., 1999. Assessment of the state of pollution by heavy metals in the superficial sediments of Algiers Bay. *Applied Radiations and Isotopes*, **50** (1999): 975-980.
- BENSAM P., 1965. On a freak embryo of the grey shark, *Carcharhinus limbatus* M. & H. *Journal of the Marine Biological Association of India*, **7** (1): 206-207.
- BLACHE J., CADENAT J., STAUCH J., 1970. Clés de détermination des poissons de mer signalés dans l'Atlantique orientale tropicale (entre le 20^{ème} parallèle N. et le 15^{ème} parallèle S). *Faune tropicale ORSTOM*, **18**: 1-479.

- BONFIL R., 1989. An abnormal embryo of the reef shark *Carcharhinus perezi* (Poey) from Yucatan, Mexico. *Northeast Gulf Science*, **10** (2): 153-155.
- BRADAI M.N., QUIGNARD J.-P., BOUAIN A., JARBOUI O., OUANNES-GHORBEL A., BEN ABDALLAH L., ZAOUALI J., BEN SALEM S., 2004. Ichtyofaune autochtone et exotique des côtes tunisiennes: Recensement et biogéographie. *Cybium*, **28** (4): 315-328.
- CAPAPÉ C., 1989. Les Sélaciens des côtes méditerranéennes: aspects généraux de leur écologie et exemples de peuplements. *Océanis*, **15** (3): 309-331.
- CAPAPÉ C., ZAOUALI J., Contribution à la biologie des Scyliorhinidæ des côtes tunisiennes. VI. *Galeus melastomus* Rafinesque, 1810. Répartition géographique et bathymétrique, sexualité, reproduction, fécondité. *Cahiers de Biologie Marine.*, **18** (4): 449-463.
- CAPAPÉ C., GUÉLORGET O., VERGNE Y., REYNAUD C., 2008 Reproductive biology of the blackmouth catshark, *Galeus melastomus* (Chondrichthyes: Scyliorhinidae) off the Languedocian coast (southern France, northern Mediterranean). *Journal of the Marine Biological Association of the United Kingdom*, **88** (2): 415-421. DOI: 10.1017/S002531540800060X.
- CAPAPÉ C., EL KAMEL-MOUTALIBI O., MNASRI N., BOUMAÏZA M., REYNAUD C., 2012. A case of hermaphroditism in Tortonese's stingray *Dasyatis tortonesei* from the Lagoon of Bizerte (northeastern Tunisia, central Mediterranean). *Acta Ichthyologica et Piscatoria*, **42** (2): 141-149. DOI: 10.3750/AIP2012.42.4.07.
- CAPAPÉ C., ALI M., SAAD A., ALKUSAIRY H., REYNAUD C., 2015a. Tail abnormalities in thornback ray *Raja clavata* (Chondrichthyes: Rajidae) from the coast of Syria (eastern Mediterranean). *Cahiers de Biologie Marine*, **56** (2):155-161.
- CAPAPÉ C., ALI M., SAAD A., REYNAUD C., 2015b. Atypical characteristics in the long-nosed skate *Dipturus oxyrinchus* (Linnaeus, 1758) from the coast of Syria (eastern Mediterranean). *Thalassia Salentina*, **37**: 71-80.
- CHAPMAN D.D., CORCORAN M.J., HARVEY G.M., MALAN S., SHIVJI M.S., 2003. Mating behavior of southern stingrays, *Dasyatis americana* (Dasyatidae). *Environmental Biology of Fishes*, **68** (3): :241-245. DOI: 10.1023/A:1027332113894.
- EHEMANN N.R., GONZÁLEZ-GONZÁLEZ L.D.V., 2018. First record of a single-clasper specimen of *Pseudobatos percellens* (Elasmobranchii: Rhinopristiformes: Rhinobatidae) from the Caribbean Sea, Venezuela. *Acta Ichthyologica et Piscatoria*, **48** (3): 261-263. DOI: 10.3750/AIEP/02341.
- FOUIAL-DJEBBAR D., DJEBBAR R., BADIJAH-HADI-AHMED A.Y., BUDZINCKI H., 2001. Level of polychlorinated biphenyls in marine environment of Algiers Bay. *Analytical Letters*, **44** (15): 2438-2456. DOI: 10.180/00032719.2011.551861.
- FRANCOUR P., BOUDOURESQUE C. F., HARMELIN J. G., HARMELIN-VIVIEN M. L., QUIGNARD J.-P., 1994. Are the Mediterranean waters becoming warmer? *Marine Pollution Bulletin*, **28** (9), 523-526.
- FROESE R., TSIKLIRAS A.C., STERGIUO K.I., 2011. Editorial note on weight-length relations of fishes. *Acta Ichthyologica et Piscatoria*, **41** (4): 261-263. DOI: 10.3750/AIP2011.41.4.01.
- GOLANI D., 2005. Check-list of the Mediterranean Fishes of Israel. *Zootaxa*, **947**: 1-200.

- HAAS D. L., EBERT D.A., 2008. First record of hermaphroditism in the Bering skate *Bathyraja interrupta*. *Northwestern Naturalist*, **89** (3): 181-185. DOI: 10.1898/NWN07-23.1.
- HEMIDA F., 2005. Les sélaciens de la côte algérienne, bisystématique des requins et des raies, écologie, reproduction et exploitation de quelques populations capturées. State Thesis, University of Sciences et Technical Subjects Houari Boumédiène, Algiers, Algeria, 272 pp.
- HEMIDA F., LADOU L., CAPAPÉ C., 2018. Impact de l'ichtyofaune non indigène de la côte algérienne: réalités et artefacts, pp. 165-169. In CIESM monograph 50 (F. Briand Ed.). Engaging marine scientists and fishers to share knowledge and perceptions – Early lessons. CIESM Publisher, Monaco and Paris, 218 pp.
- LIPEJ L., DULCIC J. 2010 - Checklist of the Adriatic Sea fishes. *Zootaxa*, **2859**: 1-92.
- LLORIS D., RUCABADO J., 1998. Guide FAO d'identification des espèces pour les besoins de la pêche. Guide d'identification des ressources marines vivantes pour le Maroc. FAO: Rome. 263 pp.
- MAURIN C., BONNET M., 1970. Poissons des côtes nord-ouest africaines (Campagnes de la «Thalassa», 1962 et 1968). *Revue des Travaux de l'Institut scientifique et technique des Pêches maritimes*, **34**: 125-170.
- PAULY D., 1983. Some simple methods for assessment of tropical fishes. *FAO Fisheries Technical Papers*, **234**: 3-10.
- QUÉRO J.-C., 1984. Scyliorhinidae. In: Fishes of the north-eastern Atlantic and the Mediterranean. Whitehead P.J.P., M.-L. Bauchot, J.-C. Hureau, J. Nielsen and E. Tortonese, (eds.), Paris, UNESCO pp. 95-100.
- QUÉRO J.-C., PORCHÉ P., VAYNE J.J. 2003 - Guide des poissons de l'Atlantique européen. Les Guides du naturaliste. Delachaux and Niestlé: Lonay (Switzerland)-Paris, 465 pp.
- QUIGNARD J. P., CAPAPÉ C. Cas d'hermaphroditisme chez *Raja miraletus* L., 1758. *Travaux du Laboratoire de biologie halieutique de, Rennes*, **6**: 133-143.
- RAFRAFI-NOUIRA, S., EL KAMEL-MOUTALIBI, O., OUNIFI-BEN AMOR K., BEN AMOR M.M., CAPAPÉ, C. A case of hermaphroditism in the common eagle ray *Myliobatis aquila* (Chondrichthyes: Myliobatidae), reported from the Tunisian coast (Central Mediterranean). *Annales, Series Historia naturalis*, **27** (1): 43-48. DOI: 10.9233 / ASHN.2017.06.
- RIBEIRO-PRADO C.C., ODDONE M.C., BUENO GONZALEZ M.M., FERREIRA DE AMORIM A., CAPAPÉ C., 2008 Morphological abnormalities in skates and rays (Chondrichthyes) from off Southeastern Brazil. *Arquivos de Ciencias do Mar, Fortaleza*, **41** (2): 21-28.
- RIBEIRO-PRADO C.C., ODDONE M.C., FERREIRA DE AMORIM A., CAPAPÉ C., 2009. An abnormal hermaphrodite pelagic stingray *Pteroplatytrygon violacea* (Dasyatidae) captured off southern coast of Brazil. *Cahiers de Biologie Marine*, **50** (1): 91-96.
- RINELLI P., BOTTARI T., FLORIO G., ROMEO T., GIORDANO D., GRECO S., 2005. Observations on distribution and biology of *Galeus melastomus* (Chondrichthyes, Scyliorhinidae) in the southern Tyrrhenian Sea (central Mediterranean). *Cybium*, **29** (1): 41-46.

- SANTANDER-NETO J., LESSA R., 2013. Hermaphroditic small-eyed roundray (*Urotrygon microphthalmum*) from north-eastern Brazil. *Marine Biodiversity Records*, **6**: e60. DOI: 10.1017/S1755267213000353.
- SERENA F., 2005 - *Field Identification Guide to the sharks and rays of the Mediterranean and Black Sea*. FAO species Identification Guide for Fisheries Purposes. FAO: Rome, 97 pp.
- SPRINGER S., 1979. A revision of the catsharks, family Scyliorhinidae. NOAA Technical Report NMFS Circular, 422, V + 152 pp.
- TORRES-HUERTA S.F., MERAZ J., CARRASCO-BAUTISTA P.E., DIAZ-CARBADILLO P.L., 2015. Morphological abnormalities of round rays of the genus *Urotrygon* in the Gulf of Tehuantepec. *Marine Biodiversity*, **46** (1): 309-315. DOI: 10.1590/S1516-89132002000200017.
- TURSI A., D'ONGHIA G., MATARRESE A., PISCITELLI G., 1993. Observations on population biology of the blackmouth catshark *Galeus melastomus* (Chondrichthyes, Scyliorhinidae) in the Ionian Sea. *Cybium*, **17** (2): 187-196.
- UNGARO N., MARANO G., MARSAN R., 1996. *Galeus melastomus* Rafinesque (Selachii, Scyliorhinidae): distribuzione e biologia sui fondi battiali del basso Adriatico. *Relazioni dell'Accademia Pugliese delle Scienze*, **49**: 195-270.
- YANO K., TANAKA S., 1989. Hermaphroditism in the lantern shark *Etmopterus unicolor* (Squalidae, Chondrichthyes). *Japanese Journal of Ichthyology*, **26** (3): 338-345. DOI: 10.1007/BF02905618.