

RESEARCH ARTICLE

Environmental heterogeneity and benthic macroinvertebrate guilds in Italian lagoons

Alberto Bassetti, Nicola Galuppo & Letizia Sabetta

Department of Biological and Environmental Sciences and Technologies
University of Salento
S.P. Lecce-Monteroni 73100 Lecce
ITALY

Abstract

- 1 - Lagoons are ecotones between freshwater, marine and terrestrial biotopes, characterized by internal ecosystem heterogeneity, due to patchy spatial and temporal distribution of biotic and abiotic components, and inter-ecosystem heterogeneity, due to the various terrestrial-freshwater and freshwater-marine interfaces.
- 2 - Here, we carried out an analysis of environmental heterogeneity and benthic macro-invertebrate guilds in a sample of 26 Italian lagoons based on literature produced over a 25 year period..
- 3 - In all, 944 taxonomic units, belonging to 13 phyla, 106 orders and 343 families, were recorded. Most species had a very restricted geographic distribution range. 75% of the macroinvertebrate taxa were observed in less than three of the twenty-six lagoons considered.
- 4 - Similarity among macroinvertebrate guilds in lagoon ecosystems was remarkably low, ranging from $10.5\% \pm 7.5\%$ to $34.2\% \pm 14.4\%$ depending on the level of taxonomic resolution.
- 5 - Taxonomic heterogeneity was due to both differences in species richness and to differences in species composition: width of seaward outlet, lagoon surface area and water salinity were the most important factors affecting species richness, together accounting for up to 75% of observed inter-lagoon heterogeneity, while distance between lagoons was the most significant factor affecting similarity of species composition.
- 6 - Both environmental filtering and passive diffusion were found to be influential processes in shaping macroinvertebrate guilds in lagoon ecosystems.

Keywords: Lagoons, macro-invertebrates, community assembly rules, niche filtering

Introduction

Community and guild structure are known to be shaped by processes occurring at very different spatial and temporal scales: from speciation events and biogeography, to small-scale biotic interactions, through diffusion processes and abiotic filtering (Zobel, 1997).

At a regional scale, historical processes shaping the species pool can be assumed to be homogeneously distributed and the relative importance of diffusion processes and niche filtering, including both abiotic and competitive filters, can be assessed. Even at this scale,

historical processes shaping the species pool can easily be identified in relation to human activities, which were particularly effective in shaping the landscape of the Mediterranean coasts. The local extinction of species and introduction of others as a consequence of human activities has shaped the regional species pool over the last 15,000 years, being spatially explicit in many cases. However, these are probably better classified as components of environmental filtering, rather than true 'historical processes'.

Transitional water ecosystems represent an ideal study environment in which to evaluate the

significance of various potential drivers of community and guild structure at the local, ecosystem, level. This is due to some features which, when combined together, make transitional waters very distinctive ecosystems in the coastal landscape. Here, we will focus on coastal lagoons, but most of the features outlined below can be generalized to other transitional water body types (Basset *et al.*, 2006; Elliott and McLusky, 2006). Coastal lagoons are habitat islands, whose boundaries and surface area can be identified in relation to tides. They are known to be harsh ecosystems for both plants and animal guilds due to their strong gradients of salinity, temperature and dissolved oxygen in space and time. High water exchange rates and heterogeneity of deposition and erosion processes are additional limiting factors for many species. Because of these two features, niche filtering can be expected to be very significant in lagoon ecosystems, selecting very similar species according to a few well known traits, of which osmoregulation (Hauxwell, *et al.*, 1998), resistance to hypoxia (Smale and Rabeni, 1995), temperature variability adaptation (Nair and Anger, 1972) are the most studied. As habitat islands, lagoons have very little endemism, if any; colonizers come directly from the surrounding freshwater and marine ecosystems, which are the source habitats of lagoons of most lagoon species, with the notable exception of many migratory birds. Since lagoons are surrounded by adequate habitats, rather than inhospitable habitats, dispersal and diffusion are expected to be relevant processes in coastal lagoons for many species and indeed entire guilds. Therefore, relative homogeneity of guild composition among lagoons could be an expected consequence of the strong filtering and dispersion influences. On the other hand, since physiographic and hydrological factors are known to differentiate lagoons into classes according to geomorphology (Kjerfve, 1994), salinity (Battaglia, 1959) and degree of confinement (Guelorget and Perthuisot, 1983; Guelorget *et al.*, 1983), strong niche filtering could also be expected to result in some degree of heterogeneity of species composition among

lagoon types. However, the relative roles of niche filtering and dispersion processes in guild and community assembly in lagoon ecosystems are still poorly investigated.

In the Mediterranean region, lagoon species and ecosystems have traditionally received much attention, mainly following either a species-centred approach, focusing on adaptations (Cognetti and Maltagliati, 2000), or a perturbation-centred approach, focusing on responses and indicators (Reizopoulou and Nicolaidou, 2004; Mouillot, *et al.*, 2006). On the other hand, only a few guild- and community-level studies have considered groups of ecosystems, searching for patterns of fish (Perez-Ruzafa, 2005) and macroinvertebrate (Reizopoulou and Nicolaidou, 2004; Sabetta *et al.*, 2007) guilds linked to environmental factors.

Here, we performed a comparative analysis of macroinvertebrate guild structure in Italian lagoons in order to make a quantitative assessment of macroinvertebrate guild heterogeneity among lagoons. Therefore, the paper assesses the contrasting hypotheses of the relative homogeneity vs. heterogeneity of lagoon guilds/communities, arising from the role of niche filtering and passive diffusion processes in structuring lagoon communities. Differences in the taxonomic structures of benthic macroinvertebrate guilds may be due to different species richness, different taxonomic composition or different species-abundance distributions. The first two components, which tend to be more conservative than the third, were directly addressed in this study.

Material and methods

Data sources

This paper is based on data-mining activity of the taxonomic composition of benthic macroinvertebrate guilds in Italian lagoons. The data used in the present study were gleaned from published sources, including international journals retrieved from electronic databases (ASFA, WebSpirs, Biological Abstracts, Current Contents and Zoological Records), Italian journals covering the fields of ecology

and zoology (*Accademia Peloritana, Il naturalista siciliano, Oebalia, Rendiconti del Seminario della Facoltà di Scienze dell'Università di Cagliari, Thalassia Salentina*) and proceedings of Italian scientific societies covering the field of aquatic ecology and zoology (*Associazione Italiana di Oceanografia e Limnologia, Società Italiana di Biologia Marina, Società Italiana di Ecologia, Società Sarda di Scienze Naturali*). The search was restricted to papers published in the last 25 years.

The electronic search was performed through a three-way factorial combination of the following groups of keywords: a) Italy, Italian; b) lagoon, coastal lake, coastal pond, saltmarsh, saltern, brackish; c) benthos, macrobenthos, macroinvertebrate, benthic invertebrates, benthic fauna. Three unpublished reports to which one of the authors directly contributed were also taken into consideration.

The dataset

In total the search produced 205 papers containing taxonomic lists of benthic macroinvertebrates in Italian lagoons; most papers focused on a single lagoon but comparative studies of different lagoons were also collected in this way. The 205 papers were screened according to three main criteria: a) taxonomic resolution (most taxa classified down to the genus or species level); b) taxonomic completeness (the taxonomic list of a lagoon, resulting from one or more papers, cannot be limited to a few selected phyla); c) sampling design (samplings carried out at a seasonal or higher resolution). According to these criteria, 169 papers were selected for the analysis, referring to 26 lagoon ecosystems. A list of the considered ecosystems, with their key structural abiotic features, is presented in Table 1. In the dataset obtained by applying these criteria, the number of articles referring to each of the 26 transitional aquatic ecosystems ranges from 1 (Massaciuccoli coastal lake, Torre Guaceto brackish wetland, Marsala saltern and Piallassa Baiona lagoon) to 22 (Venice lagoon).

For each lagoon, fifteen physiographic and hydrological parameters were collected from the

published papers as well as from published satellite images and the "Tide tables of Italy" (Istituto Idrografico della Marina, Genova 1999). The parameters are surface area, perimeter and sinuosity of the water body, maximum axis and minimum axis, length and width (and length/width ratio) of its seaward outlet(s), mean depth, maximum and minimum tide, tidal range, maximum and minimum salinity and salinity range.

Study sites

The 26 Italian coastal lagoons for which published species lists of benthic macroinvertebrate taxa were collected are distributed among 15 provinces belonging to 7 regions (Fig. 1), with 6 coastal lagoons in the region of Lazio, 5 in the region of Puglia, 4 in the regions of Sicilia and Emilia Romagna, 3 in the regions of Veneto and Toscana and 1 in the region of Friuli Venezia Giulia. In total, the 26 selected lagoons cover a coastal area of 1133.43 km², which represents 69.6% of the overall surface of Italian transitional waters.

Data analysis

The data were organized into two matrices: a presence/absence matrix with benthic macroinvertebrate data organized into 26 columns (representing the selected lagoons) and 944 rows (the macroinvertebrate taxa); and a rectangular matrix with 26 rows (representing the selected lagoons) and 17 columns, two of which describing the benthic macroinvertebrate guilds (taxonomic richness and standard body length of the largest species), nine describing the physiographic characteristics of the water bodies (surface area, perimeter, sinuosity, length of min. axis, length of max. axis, seaward outlet length, outlet width, outlet length/width ratio, average water depth), and three describing the tidal regime (i.e., minimum and maximum tides and tidal range) and the salinity regime (i.e., minimum, maximum and range of salinity).

The taxonomic composition similarity between lagoon pairs was measured using the Jaccard similarity index, as follows:

$$SJ = a/(a+b+c) \times 100$$

where

SJ = Jaccard similarity coefficient,

a = number of taxa shared by two lagoons,

b = number of taxa unique to the first lagoon,

and

c = number of taxa unique to the second lagoon.

The species area-relationships in the selected sample of Italian lagoons were analysed utilising the well-known Arrenhius power function (Arrenhius, 1921) as a descriptive model for species accumulation patterns:

$$S = cAZ, \text{ or } \log S = \log C + z \log A,$$

where A is the surface area of the lagoon, S is taxonomic richness expressed as total number of species found, and c and z are regression coefficients. Specifically, z indicates the rate of species number increasing with area and c is a fitting coefficient, or estimated number of species per "unit" area.

Multiple regression was used to evaluate the relative influence and the cumulative importance of surface area and other abiotic structural factors to macro-invertebrate taxonomic richness in the studied Italian lagoons

Results

Abiotic niche dimensions and species list

The sample of lagoons considered in this study covers most of the range of variation in Italy. The lagoon list (Table 1a) geographically includes ecosystems from both northern and southern regions, with the exception of Sardinia; in terms of surface area, tidal range, connection with the sea (outlet length and width) and water salinity, the two largest lagoon complexes, and some of the smallest lagoons, micro-tidal and non-tidal ecosystems, as well as oligohaline and polyhaline ecosystems were included. On the basis of the abiotic structural parameters considered, the 26 lagoons are clustered into 3 groups (Figure 1).

Lagoon surface, width and length of outlets and tidal range were the structural abiotic parameters which statistically differentiated the three groups (Table 1b; ordinary t-statistics, $P<0.05$); specifically, group B, including most

of the Northern Adriatic lagoons and two enclosed sea gulfs, was characterised by the widest outlets, largest surface areas and the greatest tidal ranges.

Overall, 944 taxa were reported to colonise the whole sample of lagoons, including 343 families, 106 orders, 31 classes and 13 phyla. Mollusca, Annelida and Arthropoda were the three most common phyla. A complete taxonomic list is reported as Annex 1.




Figure 1 MDS of structural abiotic similarity among lagoon ecosystems. The spatial distribution of clusters of lagoon ecosystems at a significance level $a=0.05$ is reported. Lagoons are reported with their ID number listed in Table 1

Among the Mollusca, the most common taxa in the Italian lagoons are *Cerastoderma glaucum* (Poiret, 1789), *Abra segmentum* (Récluz, 1843) and *Mytilus galloprovincialis* (Lamarck, 1819) in the Class of Bivalvia, and *Cyclope neritea* (Linné, 1758), *Bittium reticulatum* (da Costa, 1778) and *Hydrobia stagnalis* (Baster, 1765) in the Class of Gastropoda. Among the Annelida, *Polydora ciliata* (Johnston, 1838), *Capitella capitata* (Fabricius, 1780), *Ficopomatus enigmaticus* (Fauvel, 1923), *Nereis diversicolor* (O.F. Müller, 1776), *Perinereis cultrifera* (Grube, 1840) and *Malacoboceros fuliginosus* (Claparède, 1868) are the most commonly reported taxa. Finally, among the Arthropoda, the most widespread species are *Gammarus aequicauda* (Martyinov, 1931), *Corophium insidiosum* (Crawford, 1937), *Carcinus aestuarii* (Nardo, 1847), *Microdeutopus gryllotalpa* (Costa, 1853), *Gammarus insensibilis* (Stock, 1966) and *Melita palmata* (Montagu, 1804).

Table 1 – Main structural abiotic features of the 26 lagoons considered for the study (a) and of the 3 groups shown by the MDS (b). The P values of pairwise comparisons of parameter averages between groups are reported for every abiotic feature (Student's t-test)

(a)

ID	Lagoon Name	Abiotic factors								
		A	B	C	D	E	F	G	H	I
1	Grado Marano	160	2	2	485,0	2732,6	0,596	17,05	27,5	15,85
2	Venezia	549	2	1,5	1635,0	8992,5	0,576	26,3	19,4	22,5
3	Sacca Canarin	8,9	1,5	1	365,0	871,6	0,411	14,5	29	14,9
4	Scardovari	38	1,3	1,5	318,0	1011,9	0,411	18,65	29,3	14
5	Nazioni	1	2,1	2	2655,2	8,3	0,411	25	12	16,65
6	Goro	32	1,6	1,5	483,0	1705,1	0,411	21,75	26,3	16,2
7	Comacchio	115	1,2	0,8	441,1	286,7	0,411	23,1	26,2	16
8	Pialassa Baiona	11,8	1,4	0,8	1984,5	132,3	0,411	23	26	16,5
9	Massaciuccoli	7	1,6	3	4276,0	117,2	0,204	8,7	14,6	17,5
10	Orbetello	25,5	2,3	1	1838,0	92,9	0,234	34,35	15,5	17,3
11	Lesina	51	2	0,8	985,0	34,0	0,156	22,5	23	16,5
12	Varano	64	1,2	4	450,1	30,0	0,156	24,86	26,2	19,15
13	Fogliano	4	1,8	0,9	356,0	57,0	0,261	33	30	20
14	Monaci	0,95	1	0,8	818,4	27,3	0,261	27,5	21	19
15	Caprolace	2,3	1	1	974,0	23,6	0,261	41,5	17	20
16	Sabaudia	3,7	3,9	4	980,0	15,2	0,261	22,25	21,5	16,2
17	Fondi	3,9	3,9	9	2349,5	76,9	0,261	20	14	19
18	Lungo	0,47	1,3	4	213,6	17,1	0,261	20	26	20,5
19	Fusaro	1	1,2	3	630,3	39,4	0,246	37,15	8,7	18,65
20	Torre Guaceto	1,2	1,6	0,4	10,0	0,4	0,195	19	28	16
21	Acquatina	0,45	2,94	1	66,8	267,1	0,195	25	30	18
22	Alimini	1,37	2,29	1,5	146,7	19,6	0,195	29	22	19
23	Oliveri-Tindari	0,19	2,7	3	0,0	0,0	0,124	25,5	21	21,15
24	Saline di Marsala	0,89	1,4	0,6	0,0	0,0	0,156	55	30	21
25	Stagnone Marsala	24	1,8	1	314,0	1794,6	0,156	39,3	12,4	19,77
26	Rada di Augusta	23,5	1,4	14,9	235,0	1512,7	0,187	38,15	0,1	24,77

A= surface area (km²); B= sinuosity; C= mean depth (m); D= length of seaward outlet (m); E= width of seaward outlet (m); F= tidal range (m); G= mean salinity (PSU); H= range of salinity (PSU); I= mean water temperature (°C)

(b)

Group name	Parameter average								
	A	B	C	D	E	F	G	H	I
X	18,34	1,95	2,35	1197,82	77,78	0,26	26,06	20,86	18,12
Y	119,34	1,66	3,34	547,86	2660,14	0,39	25,10	20,57	18,28
Z	0,76	1,90	1,33	3,33	0,13	0,16	33,17	26,33	19,38

	P values								
	A	B	C	D	E	F	G	H	I
X,Y	0,005	0,573	0,766	0,280	0,000	0,025	0,750	0,363	0,899
X,Z	0,126	0,964	0,336	0,000	0,000	0,044	0,440	0,215	0,307
Y,Z	0,003	0,487	0,414	0,000	0,000	0,047	0,446	0,461	0,632

Heterogeneity of macroinvertebrate guilds in lagoon ecosystems

Most of the macroinvertebrate taxa which were found to colonise the sample of 26 Italian lagoons were actually reported for a very restricted number of lagoons. 60.6% of taxa were reported for a single lagoon, and only 5.9% of taxa (i.e., 56 taxa) were recorded for at least 8 lagoons (see Annex 1). These most widespread taxa included 20 species of Annelida, 18 species of Arthropoda and 15 species of Mollusca. In the data-base utilised for

the study, the number of taxa recorded in a fixed number of lagoons was inversely related to the number of lagoons (Figure 2).

Species richness varied by more than one order of magnitude between lagoons. Venice lagoon and the Stagnone di Marsala had the highest taxonomic richness at every level of taxonomic resolution considered in the study (S=285 and S=220 at the lowest level of taxonomic resolution, respectively). On the other hand, Lago Nazioni and Torre Guaceto had the lowest species richness at every level of taxonomic

resolution considered in the study ($S=21$ and $S=22$ at the lowest level of taxonomic resolution, respectively).




Figure 2 – Width of the distribution of benthic macroinvertebrates within the considered sample of Italian lagoons, as expressed by the number of lagoons colonised by the 944 taxa.

Average similarity in the taxonomic composition of benthic macroinvertebrate guilds was low between lagoons at every level of taxonomic resolution considered, being 0.105 ± 0.008 at the taxon level, 0.217 ± 0.013 at the family level and 0.346 ± 0.016 at the order level. Average similarity between lagoon pairs ranged from 0 to 0.345 at the taxon level, being highest for the spatially neighbouring Fogliano and Monaci lagoons; it ranged from 0 to 0.55 at the family level, being highest for the spatially neighbouring Goro and Sacca del Canarin lagoons, and from 0 to 0.69 at the order level, being highest again for the Goro and Sacca del Canarin lagoons. Taxonomic similarity linearly increased when species composition was progressively restricted to species/taxa with increasingly wider distribution (i.e., removing progressively the species/taxa occurring in an increasingly higher number of lagoons, from 1 to eighteen) (Figure 3a). However, even when almost 95% of species were removed, considering only those species occurring in at least 8 lagoon ecosystems, average similarity in the taxonomic composition of benthic macroinvertebrates among lagoons was still lower than 30% (Figure 3a; $J=0.26 \pm 0.014$). The removal of the rarest species had the most significant effect on taxonomic similarity among lagoons (Figure 3b); however, highly

significant effects were also observed at the thresholds of species occurring in at least 10, 12 and 15 lagoons (black dots in Figure 3b).

Determinants of macro-invertebrate guild heterogeneity

A significant component of the variability of both species richness and taxonomic similarity among lagoons was found to be related to some key physiographic, hydrological and physical-chemical factors.

Outlet width and surface area of coastal lagoons were the only two structural factors considered in the study to which species richness was directly related. The species-area relationship had a slope of 0.18 and explained more than 25% of variability in species richness among the considered lagoons (Figure 4). At the multivariate level, the integration of either outlet width or surface area with average water salinity explained respectively 70% and 48% of species richness variability among lagoons.




Figure 3 – (a) Relationship between taxonomic similarity and number of taxa. Error bars represent \pm SD of mean. (b) Relationship between significance of taxonomic similarity variation and number of taxa removed.

Distance between lagoons and average water salinity were the two factors with which taxonomic dissimilarity of macroinvertebrate species

composition at every level of taxonomic resolution and at every threshold of species spatial niche (as quantified by the number of lagoons colonised) showed the strongest and most consistent relationships (Figure 5). At the multivariate level, four factors (distance between lagoons, surface area, water salinity and water depth) were the most important factors affecting dissimilarity between taxonomic composition of benthic

macroinvertebrates at every level or condition considered (Table 2). When a threshold is imposed on the spatial niche of a species, hydrodynamic factors, such as outlet length and width and tidal range also had statistical relevance (Table 2). The variance explained by the significant factors listed in Table 2 range from 21 %, when order level was considered, to 44 %, when taxa occurring in at least 15 lagoons were considered.

Table 2 Multiple regression analysis of the relationship between structural abiotic characteristics of the considered lagoons and taxonomic composition of benthic macro-invertebrate guilds. Dissimilarity between lagoons in the two sets of data were compared for the different levels of taxonomic resolution and for the taxa occurring in a number of lagoons larger than some threshold level (taxa=all taxa considered; T12=only taxa occurring in at least 12 lagoons; T15= only taxa occurring in at least 15 lagoons). Beta values and significance levels are also shown (Stepwise multiple regression; ***= P<0.001; **= P<0.01).

	1	2	3	4	5	6	7
taxa	0,44 ***	0,22 ***	-0,2 ***	0,11 **			
family	0,32 ***	0,18 ***	-0,25 ***	0,17 ***			
order	0,23 ***	0,17 **	-0,19 ***	0,15 **			
T12	0,44 ***		-0,29 ***	0,17 **	-0,25 ***		
T15	0,59 ***	0,1 **	-0,23 ***	0,11 **	-0,32 ***	0,14 **	-0,25 ***

1= Distance (km); 2=Mean Salinity (PSU); 3=Surface (Km²); 4=Mean Depth (m); 5=Length out. (m); 6=Width out. (m); 7=Tidal range (m)

Discussion

A central aim of community ecology has been, and remains, to decode patterns of species abundance into mechanisms of guild and community assembly. Here, using lagoons as model ecosystems, we have searched for patterns of species abundance and species association in a sample of Italian lagoons, looking for the influence of niche filtering as a mechanism underpinning community organization. The 26 considered lagoons were likely to be a representative sample of Italian lagoons, accounting for 14.8% of lagoon numbers and 69.6% of lagoon surface area in Italy (Basset and Sabetta, 2005).

The results of this study completely rejected any hypothesis of relative homogeneity of the taxonomic composition of macroinvertebrate guilds in lagoon ecosystems, either due to strong and directional niche filtering and/or to passive diffusion of colonising species.

Taxonomic composition of macroinvertebrate guilds was extremely heterogeneous among lagoons at every level of taxonomic resolution and selection considered.




Figure 4 Species-area relationships in lagoon ecosystems.

This was highlighted at the species level, since almost 95% of the 944 species detected overall were recorded in less than 8 lagoons, and at the guild level, as shown by the low similarity between lagoons. Even neighbouring lagoons of similar surface area, which would be expected

to share a common potential species pool, such as Lesina and Varano in the Puglia region or Caprolace and Monaci in the Lazio region, had very different taxonomic composition of macroinvertebrate guilds (Jaccard similarity index; $J = 0.174$ and $J = 0.154$, respectively). In principle, the diversity of macro-invertebrate guilds could result from stochastic, ecological and evolutionary processes on the local (e.g.,

community level) and regional (e.g., meta-community level) scale (Hubbell, 2001), or from dispersal and recruitment limitations on macro-invertebrate taxa (Hurt & Pacala, 1995), or from the structural heterogeneity of the transitional waters (Basset & Abbiati, 2004) and the consequent selection of macroinvertebrate taxa according to their functional traits and niche requirements (MacArthur, 1970).




Figure 5 Relationships between taxonomic dissimilarity and distance dissimilarity / salinity dissimilarity for taxa occurring in a number of lagoons considered (a, b) and at each of the different levels of taxonomic resolution (c, d, e).

The results of this study showed that in Mediterranean lagoons, niche filtering of species traits along the main axes of structural heterogeneity of lagoon ecosystems has an important role in the assembly of macroinvertebrate guilds. However, the heterogeneity of macro-invertebrate guilds among the Italian lagoons is much larger than that explained by niche filtering. Therefore, the observed results suggest that other processes, either stochastic, such as diffusion and colonisation, or deterministic, such as competitive selection, or an integration of stochastic and deterministic processes in a competitive lottery must have an important effect on niche filtering in the assembly of benthic macro-invertebrate guilds.

The deterministic influence of some niche dimensions of lagoon ecosystems, such as surface area, water salinity, outlet width and length and water depth, on taxonomic abundance and taxa distribution among lagoons of benthic macro-invertebrates supports the role of niche filtering. Within and among habitats, structural heterogeneity is a common feature of lagoons and coastal lakes, which are ecotones between terrestrial, freshwater and marine ecosystems (e.g., Basset and Abbiati, 2004), whose relative influence varies over time and determines characteristic scales of temporal variability inside transitional aquatic ecosystems (Comin *et al.*, 2004). The observation of significant co-variations of taxonomic richness and taxonomic composition of benthic macro-invertebrates along gradients of surface area, water salinity, outlet width and length and water depth, agree with various classifications already proposed to account for the heterogeneity of transitional Mediterranean waters, based on geomorphology (Kjerfve, 1994), salinity (Battaglia, 1959), degree of confinement (Guelorget and Perthuisot, 1983; Guelorget *et al.*, 1983) and physiographic characteristics, such as surface area and outlet length (Basset *et al.*, 2006). Physiographic characteristics have recently been proposed as major factors determining lagoon type across a very large geographical range from the Mediterranean (Basset *et al.*, 2005) to French

Polyynesia (Andrefouet *et al.*, 2001). Consequently, surface area, water salinity, and outlet width and length, which are likely to be descriptors of water exchange and confinement of the lagoon ecosystems, can actually be considered the key limiting, dimensions defining the environmental niche space for benthic macroinvertebrates in lagoon ecosystems. Physiographic and hydrological differences among lagoons set different environmental niche conditions, within which there is ground for niche partitioning among species. Niche differences among species with respect to the physical environment could potentially account for the heterogeneity observed in the taxonomic composition of macro-invertebrate guilds in the 26 Italian lagoons considered, particularly as regards species richness. It is supported by the observation of significant species/area relationships, which have already been demonstrated for lagoon ecosystems (Sabetta *et al.*, 2007). Species/area relationships in lagoon ecosystems are held to be consistent with a deterministic integration of the capacity rule (*sensu* Brown, 1981) set by energy availability, according to the More Individual Hypothesis (MIH, Wright, 1983), and the allocation rule arising from species interactions and niche partitioning (Sabetta *et al.*, 2007).

On the other hand, while up to 75% of taxonomic richness variability among Italian lagoons was explained by outlet width, surface area and average salinity of Italian lagoons, only 44% of the heterogeneity in taxonomic composition was explained. Moreover, distance between lagoons was the most important source of variation of taxonomic composition, even though the above-mentioned structural niche dimensions were also significant sources of variation. Observations of exponential decline in taxonomic similarity with increasing geographic distance are common for many different guilds, including fossil plant communities (Enquist *et al.*, 2002), different terrestrial guilds of the Canary Islands (Fernandes-Palacios and Andersson, 1993) and aquatic parasites (Poulin, 2003); in all cases, dispersal was considered a main factor

determining the observed patterns of taxonomic similarity decline with increasing geographic distance. Therefore, dispersal is likely to be influential in the assembly of macroinvertebrate guilds in the studied Italian lagoons. However, the patterns of taxonomic composition similarity among lagoons do not appear to be accounted for completely by potential limitations on the dispersal and recruitment of macro-invertebrate taxa. Indeed, even though neighbouring lagoons, which would be expected to share common hydrology and recruitment from marine and freshwater environments, tend to have more similar taxonomic composition than lagoons which are geographically distant, this similarity is in any case very low, even when comparing neighbouring lagoons of similar surface area, such as Lesina and Varano or Caprolace and Monaci. Dispersal and recruitment limitation could explain the meta-community level heterogeneity and could have a role in the geographical clustering of Italian lagoons (Sabetta *et al.*, 2004), but they do not seem to account for the variations observed at a smaller spatial geographical scale. Therefore, the observed heterogeneity of macroinvertebrate guilds supports the view that some complementarities among dispersal and recruitment, differences in the freshwater and marine fluxes of both abiotic components among lagoons, and directional or neutral (He, 2005) species interactions have determined over time such a large gradient of environmental conditions that each lagoon has its own specific features, implying an individualistic approach to lagoon ecology.

In conclusion, the observed results exclude any hypothesis of large scale homogeneity of macro-invertebrate guilds of Mediterranean lagoons, due to expected strong filtering selection and dispersal. Moreover, both niche filtering and dispersal and recruitment were found to have an important role in the assembly of macro-invertebrate guilds, even though the presence of some unexplained variations suggest species interaction at the local level.

References

- Andréfouët S, Claereboudt M, Matsakis P, Pagès J, Dufour P 2001. Typology of atoll rims in Tuamotu Archipelago (French Polynesia) at landscape scale using SPOT HRV images. International Journal of Remote Sensing 22(6): 987-1004 (18).
- Arrhenius O 1921. Species and area. Journal of Ecology 9: 95-99.
- Basset A 1994. Mole corporea ed organizzazione delle guild a base detrito all'interfaccia terra-acqua dolce. Atti Congresso S.I.T.E. 16: 155-160.
- Basset A 1995. Body size-related coexistence: an approach through allometric constraints on home-range use. Ecology 76: 1027-1035.
- Basset A, Abbiati M 2004. Challenges to transitional water monitoring: ecological descriptors and scales. Aquatic Conservation: Marine and Freshwater Ecosystems 14: S1-S3.
- Basset A, Sabetta L 2005. Identificazione e caratterizzazione degli ambienti acquatici di transizione: risultati di una indagine italiana a livello nazionale. Atti Workshop "Il monitoraggio delle acque di transizione". Venezia 27-29 ottobre 2004: 78-86.
- Basset A, Sabetta L, Fonnesu A, Mouillot D, Do Chi T, Viaroli P, Giordani G, Reizopoulou S, Abbiati M, Carrada GC 2006. Typology in Mediterranean transitional waters: new challenges and perspectives. Aquatic Conservation: Marine and Freshwater Ecosystems 16(5): 441 - 455.
- Battaglia B 1959. Final resolution of the symposium on the classification of brackish waters. Archivio Di Oceanografia E Limnologia 11(suppl): 243-248.
- Bonn A, Storch D, Gaston KJ 2004. Structure of the species-energy relationship. Proceeding: Biological Sciences 271: 1685-1691.
- Brown JH 1981. Two decades of homage to Santa Rosalia: toward a general theory of diversity. American Zoologist 21: 877-888.
- Carrada G.C., Cicogna F., Fresi E. 1988. Le lagune costiere: ricerca e gestione. CLEM Pubbl, Napoli, 254 pp.
- Comin FA, Menendez M, Herrera JA 2004. Spatial and temporal scales for monitoring coastal aquatic ecosystems. Aquatic Conservation: Marine and Freshwater Ecosystems 14: S5-S17.
- Cognetti G 1982. Adaptive strategy of brackish-water fauna in pure and polluted waters. Marine Pollution Bulletin 13: 247-250.
- Cognetti G, Maltagliati F 2000. Biodiversity and adaptive mechanisms in brackish water fauna. Marine Pollution Bulletin, 40: 7-14.
- Connor EF, McCoy ED 1979. The statistics and biology of the species-area relationship. The American Naturalist 113: 791-833.

- Enquist BJ, Haskell JP, Tiffney BH 2002. General patterns of taxonomic and biomass partitioning in extant and fossil plant communities. *Nature* 419: 610-613.
- Evans KL, Greenwood JJD, Gaston KJ 2005. The role of extinction and colonisation in generating species-energy relationships. *Journal of Animal Ecology* 74: 498-507.
- Fernandez-Palacios JM, Andersson C 1993. Species composition and within archipelago co-occurrence patterns in the Canary Islands. *Ecography* 16: 31-36.
- Guelorget O, Perthuisot JP 1983. Le domaine paralique. Expressions géologiques, biologiques et économiques du confinement. *Travaux du laboratoire de géologie* 16: 1-136.
- Guelorget O, Frisoni GF, Perthuisot JP 1983. Zonation biologique des milieux lagunaires: définition d'une échelle de confinement dans le domaine paralique méditerranéen. *Journal de Recherche océanographique* 8: 15-35.
- Hauxwell J, McClelland J, Behr PJ, Valiela I 1998. Relative importance of grazing and nutrient controls of macroalgal biomass in three temperate shallow estuaries. *Estuaries* 21(2): 347 - 360.
- Hawkins BA, Field R, Cornell HV, Currie DJ, Guegan JF, Kaufman DM, Kerr JT, Mittelbach GG, Oberdorff T, O'Brien EM, Porter EM, Turner JRG 2003. Energy, water and broad scale patterns of species richness. *Ecology* 84: 3105- 3117.
- He F 2005. Deriving a neutral model of species abundance from fundamental mechanisms of population dynamics. *Functional Ecology* 19: 187-193.
- Holt RD, Lawton JH, Polis GA, Martinez ND 1999. Trophic rank and the species-area relationship. *Ecology* 80: 1495-1504.
- Hubbell SP 2001. The unified neutral theory of biodiversity and biogeography. Princeton University Press, Princeton, 448 pp.
- Hurt GC, Pacala SW 1995. The consequences of recruitment limitation. Reconciling chance, history and competitive differences between plants. *Journal of Theoretical Biology* 176: 1-12.
- Istituto Idrografico della Marina, 1999. Tavola italiana delle maree, Genova, 118 pp.
- Kjerfve B. 1994. Coastal Lagoon Processes. Elsevier, Amsterdam, 577 pp.
- MacArthur RH 1970. Species packing and competitive equilibrium for many species. *Theoretical Population Biology* 1: 1-11.
- MacArthur R.H. 1972. Geographical Ecology. Harper & Row, N.Y., 269 pp.
- MacArthur R.H., Wilson E.O. 1967. The Theory of Island Biogeography. Princeton Univ. Press, Princeton, N.J., 203 pp.
- McLusky DS, Elliott M 2007. Transitional waters: A new approach, semantics or just muddying the waters? *Estuarine, Coastal and Shelf Science* 71: 359-363.
- Moullot D, Spatharis S, Reizopoulou S, Laugier T, Sabetta L, Bassett A., Do Chi T 2006. Alternatives to taxonomic-based approaches to assess changes in transitional water communities. *Aquatic Conservation: Marine And Freshwater Ecosystems* 16: 469-482.
- Nair KKC, Anger K 1979. Life cycle of *Corophium insidiosum* (Crustacea, Amphipoda) in laboratory culture. *Helgol. Wiss. Meeresunters* 32: 279 - 294.
- Pérez-Ruzafa A, Quispe-Becerra JI, García-Charton JA, Marcos C 2004. Composition, structure and distribution of the ichthyoplankton in a Mediterranean coastal lagoon. *Journal of Fish Biology* 64(1): 202 – 218.
- Perez-Ruzafa A, Fernandez AI, Marcos C, Gilabert J, Quispe JI, Garcia-Charton JA 2005. Spatial and temporal variations of hydrological conditions, nutrients and chlorophyll a in a Mediterranean coastal lagoon (Mar Menor, Spain). *Hydrobiologia* 550:11-27
- Peters R.H. 1983. The Ecological Implications of Body Size. Cambridge University Press, Cambridge, 344 pp.
- Poulin R 2003. The decay of similarity with geographical distance in parasite communities of vertebrate hosts. *Journal of Biogeography* 30: 1609-1615
- Reizopoulou S, Nicolaidou A 2004. Benthic diversity of coastal brackish-water lagoons in western Greece. *Aquatic Conservation: Marine And Freshwater Ecosystems* 14: S93-S102
- Sabetta L, Fiocca A, Margheriti L, Vignes F, Bassett A, Mangoni O, Carrada GC, Ruggieri N, Ianni C 2004. Phytoplankton size structure and environmental forcing within the euphotic zone in the southern adriatic-ionian coastal area. *Chemistry and Ecology* 20(1): S303-S317.
- Sabetta L, Barbone E, Giardino A, Galuppo N, Bassett A 2007. Species-area patterns of benthic macro-invertebrates in Italian lagoons. *Hydrobiologia* 577: 127-139
- Smale MA, Rabeni CF 1995. Hypoxia and hyperthermia tolerances of headwater stream fishes. *Transactions of the American Fisheries Society* 124(5): 698-710.
- WFD. 2000. European Commission Directive 2000/60/EC of the European Parliament and of the Council of 23 October 2000 establishing a framework for Community action in the field of water policy. Official Journal of the European Communities, L 327/1-72, Brussels.
- Williams C.B. 1964. Patterns in the Balance of Nature. Academic Press, London, 324 pp.

- Wright DH 1983. Species-energy theory: an extension of species area theory. *Oikos* 41: 496–506.
- Zobel M 1997. The relative role of species pools in determining plant species richness: an alternative explanation of species coexistence? *Trends in Ecology and Evolution* 12(7): 266-269

Annex 1

Phylum	Taxon	N. sites
Mollusca	<i>Cerastoderma glaucum</i>	21
Anellida	<i>Polydora ciliata</i>	19
Mollusca	<i>Abra segmentum</i>	18
Arthropoda	<i>Gammarus aequicauda</i>	18
Arthropoda	<i>Corophium insidiosum</i>	18
Anellida	<i>Canthella capitata</i>	17
Arthropoda	<i>Carcinus estuarii</i>	16
Anellida	<i>Ficopomatus enigmaticus</i>	15
Mollusca	<i>Cyclospira nerita</i>	15
Arthropoda	<i>Microdeutonous cylindralpa</i>	14
Anellida	<i>Nereis diversicolor</i>	14
Arthropoda	<i>Gammarus insensibilis</i>	14
Arthropoda	<i>Melita palmata</i>	14
Mollusca	<i>Mytilus galloprovincialis</i>	13
Arthropoda	<i>Balanus amphitrite</i>	12
Mollusca	<i>Loripes lacteus</i>	12
Mollusca	<i>Nassarius reticulatus</i>	12
Arthropoda	<i>Balanus ebureus</i>	12
Anellida	<i>Perimereis cultirifera</i>	12
Anellida	<i>Malacoboceras fuliginosus</i>	12
Anellida	<i>Neanthes caudata</i>	11
Mollusca	<i>Bittium reticulatum</i>	11
Arthropoda	<i>Lekanesphaera hookeri</i>	11
Anellida	<i>Neanthes succinea</i>	11
Anellida	<i>Platynereis dumerelii</i>	11
Anellida	<i>Cirriformia tentaculata</i>	11
Mollusca	<i>Tapes decussatus</i>	11
Anellida	<i>Nereiphiyla rubiginosa</i>	10
Arthropoda	<i>Brachynotus sexdentatus</i>	10
Bryozoa	<i>Conopeum seurati</i>	10
Anellida	<i>Spiro decoratus</i>	10
Anellida	<i>Syllis gracilis</i>	10
Mollusca	<i>Mytilaster minimus</i>	10
Arthropoda	<i>Sphaeroma serratum</i>	9
Anellida	<i>Heteromastus filiformis</i>	9
Arthropoda	<i>Balanus improvisus</i>	9
Arthropoda	<i>Idotea balthica</i>	9
Arthropoda	<i>Chironomus salinarius</i>	9
Arthropoda	<i>Dexamine spinosa</i>	9
Arthropoda	<i>Palaeomon adspersus</i>	9
Cnidaria	<i>Ventromma haleciooides</i>	9
Mollusca	<i>Venerupis aurea</i>	9
Mollusca	<i>Hydrobia stagnalis</i>	8
Mollusca	<i>Cerithium vulgatum</i>	8
Mollusca	<i>Nassarius corniculus</i>	8
Arthropoda	<i>Corophium acherusicum</i>	8
Arthropoda	<i>Cyathura carinata</i>	8
Anellida	<i>Hydrodromes dianthus</i>	8
Anellida	<i>Naineris laevigata</i>	8
Anellida	<i>Nephrys hombergii</i>	8
Anellida	<i>Pseudopolydora antennata</i>	8
Mollusca	<i>Haminoea navicularia</i>	8
Mollusca	<i>Hydrobia sp.</i>	8
Cnidaria	<i>Obelia dichotoma</i>	8
Anellida	<i>Hydrodromes elegans</i>	8
Anellida	<i>Podarke pallida</i>	8
Arthropoda	<i>Corophium orientale</i>	7
Mollusca	<i>Haminoea hyalatis</i>	7
Arthropoda	<i>Gammarus sp.</i>	7
Cnidaria	<i>Paramemonea cinerea</i>	7
Anellida	<i>Streblospio shrusolii</i>	7
Anellida	<i>Armandia cirrhosa</i>	7
Mollusca	<i>Chamelea gallina</i>	7
Anellida	<i>Pomatoceros lamarcii</i>	7
Bryozoa	<i>Bowerbankia gracilis</i>	7
Cnidaria	<i>Laomedea angulata</i>	7
Arthropoda	<i>Palaemon elegans</i>	7
Arthropoda	<i>Caprella equilibra</i>	7
Arthropoda	<i>Tanais dulongii</i>	7
Anellida	<i>Spiro filicornis</i>	7
Mollusca	<i>Ostrea edulis</i>	6
Mollusca	<i>Tapes philippinarum</i>	6
Echinodermata	<i>Amphipholis squamata</i>	6
Anellida	<i>Nereis folsa</i>	6
Arthropoda	<i>Palaemon serratus</i>	6
Mollusca	<i>Parvicardium exiguum</i>	6
Arthropoda	<i>Upogebia pusilla</i>	6
Bryozoa	<i>Bugula stolonifera</i>	6
Mollusca	<i>Gastrana fragilis</i>	6
Mollusca	<i>Hexaplex trunculus</i>	6
Arthropoda	<i>Lekanesphaera monodi</i>	6
Anellida	<i>Marpphysa sanguinea</i>	6
Cnidaria	<i>Actinia sp.</i>	6
Echinodermata	<i>Asterina gibbosa</i>	6
Bryozoa	<i>Bugula neritina</i>	6
Anellida	<i>Prionospio cirrifera</i>	6
Anellida	<i>Terebella lapidaria</i>	6
Bryozoa	<i>Victoriella pavida</i>	6
Arthropoda	<i>Corophium acutum</i>	6

Arthropoda	<i>Erichtonius brasiliensis</i>	6
Anellida	<i>Glycera convoluta</i>	6
Anellida	<i>Prionospio multibranchiata</i>	6
Anellida	<i>Mysta picta</i>	5
Mollusca	<i>Corbula gibba</i>	5
Anellida	<i>Notomastus latericeus</i>	5
Mollusca	<i>Gibbula adriatica</i>	5
Tunicata	<i>Botryllus schlosseri</i>	5
Arthropoda	<i>Crangon crangon</i>	5
Mollusca	<i>Crassostrea gigas</i>	5
Mollusca	<i>Gibbula adansonii</i>	5
Arthropoda	<i>Idotea chelifipes</i>	5
Anellida	<i>Lumbibrineris latreilli</i>	5
Cnidaria	<i>Obelia bidentata</i>	5
Anellida	<i>Pileolaria pseudomilitaris</i>	5
Bryozoa	<i>Schizoporella errata</i>	5
Bryozoa	<i>Cryptosula pallasiiana</i>	5
Arthropoda	<i>Diogenes pugilator</i>	5
Anellida	<i>Owenta fusiformis</i>	5
Mollusca	<i>Paphia aerea</i>	5
Mollusca	<i>Mytilaster marioni</i>	5
Arthropoda	<i>Leptocheirus pilosus</i>	5
Bryozoa	<i>Zoobotryon verticillatum</i>	5
Anellida	<i>Pectinaria Koreni</i>	4
Mollusca	<i>Cerastoderma edule</i>	4
Mollusca	<i>Anadara inaequivalvis</i>	4
Cnidaria	<i>Cordylophora caspia</i>	4
Arthropoda	<i>Leptocheira savignyi</i>	4
Mollusca	<i>Muscularia senhousia</i>	4
Mollusca	<i>Venerupis corrugata</i>	4
Mollusca	<i>Cerithium rupstre</i>	4
Mollusca	<i>Nassarius mutabilis</i>	4
Mollusca	<i>Abra alba</i>	4
Arthropoda	<i>Amphiothe ramondi</i>	4
Arthropoda	<i>Diamysis bahirensis</i>	4
Mollusca	<i>Lentidium mediterraneum</i>	4
Arthropoda	<i>Orchestia mediterranea</i>	4
Endoprocta	<i>Barentia benedeni</i>	4
Mollusca	<i>Bulla striata</i>	4
Anellida	<i>Eunice vittata</i>	4
Mollusca	<i>Spisula subtruncata</i>	4
Anellida	<i>Janua pseudocorrugata</i>	4
Mollusca	<i>Mitilaster lineatus</i>	4
Anellida	<i>Nereis sp.</i>	4
Anellida	<i>Phyllocladus (Anaitides) mucosa</i>	4
Arthropoda	<i>Pilumnus hirtellus</i>	4
Anellida	<i>Aonides oxycarpa</i>	4
Endoprocta	<i>Barentia gracilis</i>	4
Arthropoda	<i>Elasmopus ranax</i>	4
Arthropoda	<i>Melita hergensis</i>	4
Anellida	<i>Prionospio malnigreni</i>	4
Mollusca	<i>Scrobicularia plana</i>	4
Anellida	<i>Syrveta plicata</i>	4
Anellida	<i>Harmothoe sp.</i>	4
Mollusca	<i>Spirilla neapolitana</i>	3
Anellida	<i>Syllis vitata</i>	3
Anellida	<i>Syllis proliferata</i>	3
Mollusca	<i>Crassostrea sp.</i>	3
Anellida	<i>Polycirrus sp.</i>	3
Arthropoda	<i>Caprella acanthifera</i>	3
Mollusca	<i>Conus mediterraneus</i>	3
Arthropoda	<i>Echinogammarus planicrurus</i>	3
Anellida	<i>Pomatoceros triquetus</i>	3
Arthropoda	<i>Pseudoleptochelia anomala</i>	3
Anellida	<i>Aphelochaeta marioni</i>	3
Mollusca	<i>Gibbula ardens</i>	3
Mollusca	<i>Nucula nucleus</i>	3
Mollusca	<i>Rissoa ventricosa</i>	3
Anellida	<i>Syllis armillaris</i>	3
Anellida	<i>Amanthia lendigera</i>	3
Arthropoda	<i>Hexapleomera robusta</i>	3
Anellida	<i>Schistomeris rudolphi</i>	3
Anellida	<i>Eudalia viridis</i>	3
Porifera	<i>Halichondria panicea</i>	3
Mollusca	<i>Macraea stultorum</i>	3
Echinodermata	<i>Paracentrotus lividus</i>	3
Anellida	<i>Phyllocladus lineata</i>	3
Bryozoa	<i>Scrupocellaria bertholletii</i>	3
Mollusca	<i>Tellina distorta</i>	3
Mollusca	<i>Abra prismatica</i>	3
Arthropoda	<i>Balanus sp.</i>	3
Anellida	<i>Janua pagensechieri</i>	3
Arthropoda	<i>Leucocthea spinicarpa</i>	3
Mollusca	<i>Modiolus barbatus</i>	3

Tunicata	<i>Molgula sp.</i>	3
Bryozoa	<i>Noellea stipata</i>	3
Echinodermata	<i>Ophiothrix fragilis</i>	3
Arthropoda	<i>Pachygrapsus marmoratus</i>	3
Arthropoda	<i>Periculodes aequimanus</i>	3
Mollusca	<i>Physella acuta</i>	3
Mollusca	<i>Retusa mammillata</i>	3
Anellida	<i>Serpula vermicularis</i>	3
Mollusca	<i>Sriarca lactea</i>	3
Anellida	<i>Syllis variegata</i>	3
Mollusca	<i>Tellina sp.</i>	3
Arthropoda	<i>Chironomus plumosus</i>	3
Arthropoda	<i>Corophium sp.</i>	3
Arthropoda	<i>Elasmosopus prociliimanus</i>	3
Bryozoa	<i>Electra monostachys</i>	3
Arthropoda	<i>Erichtonius difformis</i>	3
Anellida	<i>Eteone picta</i>	3
Mollusca	<i>Gibbula varia</i>	3
Anellida	<i>Hydrodoides pseudocircinata</i>	3
Arthropoda	<i>Jassa marmorata</i>	3
Anellida	<i>Lanice conchilega</i>	3
Anellida	<i>Lumbibrineris impatiens</i>	3
Arthropoda	<i>Paracerceis sculpta</i>	3
Anellida	<i>Paridoneis lyra</i>	3
Anellida	<i>Prionospio caspersi</i>	3
Arthropoda	<i>Pseudonotropela phasma</i>	3
Mollusca	<i>Rissoa sp.</i>	3
Bryozoa	<i>Schizoporella unicornis</i>	3
Anellida	<i>Serpula concharum</i>	3
Anellida	<i>Syllis hyalina</i>	3
Anellida	<i>Vermiliopsis striaticeps</i>	3
Cnidaria	<i>Obelia geniculata</i>	3
Mollusca	<i>Gammarellus fucicola</i>	2
Mollusca	<i>Nassarius cuvierii</i>	2
Anellida	<i>Scolelepis cirratus</i>	2
Mollusca	<i>Macoma tenuis</i>	2
Arthropoda	<i>Colomastix pusilla</i>	2
Porifera	<i>Halicondria bowerbanki</i>	2
Anellida	<i>Pholoe minuta</i>	2
Anellida	<i>Podarceopsis capensis</i>	2
Arthropoda	<i>Upogebia tipica</i>	2
Mollusca	<i>Ovatella myosotis</i>	2
Cnidaria	<i>Bougainvillia muscus</i>	2
Arthropoda	<i>Hyale crassipes</i>	2
Mollusca	<i>Lucinella divaricata</i>	2
Anellida	<i>Paleanous debilis</i>	2
Bryozoa	<i>Bugula falva</i>	2
Porifera	<i>Hymeniacidon sanguinea</i>	2
Mollusca	<i>Iris irus</i>	2
Anellida	<i>Phylodoce sp.</i>	2
Anellida	<i>Polyvithamus pictus</i>	2
Arthropoda	<i>Apseudes latreillei</i>	2
Bryozoa	<i>Bulbella abscondita</i>	2
Cnidaria	<i>Cereus pedunculatus</i>	2
Arthropoda	<i>Eriphia verrucosa</i>	2
Arthropoda	<i>Hyale perieri</i>	2
Anellida	<i>Melima palmata</i>	2
Mollusca	<i>Smaragdia viridis</i>	2
Bryozoa	<i>Tricellaria inopinata</i>	2
Anellida	<i>Syllis krohni</i>	2
Mollusca	<i>Anomia ephippium</i>	2
Mollusca	<i>Bursatella leachii</i>	2
Arthropoda	<i>Euraphia depressa</i>	2
Arthropoda	<i>Echinogammarus olivii</i>	2
Mollusca	<i>Glaes trapezia</i>	2
Mollusca	<i>Lepidothrix cinerea</i>	2
Anellida	<i>Notomastus sp.</i>	2
Arthropoda	<i>Penaeus kerathurus</i>	2
Mollusca	<i>Rapania venosa</i>	2
Arthropoda	<i>Stenothoe monoculoides</i>	2
Mollusca	<i>Tricolla pullus</i>	2
Cnidaria	<i>Aiptasia diaphana</i>	2
Cnidaria	<i>Anemonia sulcata</i>	2
Mollusca	<i>Buccinulum corneum</i>	2
Anellida	<i>Capitomastus minimus</i>	2
Arthropoda	<i>Corophium volutator</i>	2
Mollusca	<i>Donax venustus</i>	2
Anellida	<i>Eteone sp.</i>	2
Anellida	<i>Malmgreniella lunulata</i>	2
Anellida	<i>Hydrodoides sp.</i>	2
Cnidaria	<i>Laomedea calciferula</i>	2
Anellida	<i>Nereis pelagica</i>	2
Anellida	<i>Salvatoria clavata</i>	2
Mollusca	<i>Rissoa lineolata</i>	2
Anellida	<i>Sabellaria spinulosa</i>	2
Porifera	<i>Tedania anhelans</i>	2
Mollusca	<i>Theodoxus fluviatilis</i>	2
Mollusca	<i>Tricolla tenuis</i>	2
Anellida	<i>Trypanosyllis zebra</i>	2

Phylum	Taxon	N. sites
Cnidaria	<i>Tubularia crocea</i>	2
Anellida	<i>Solis torquata</i>	2
Porifera	<i>Halticola sp.</i>	2
Bryozoa	<i>Aetea truncata</i>	2
Anellida	<i>Amphipelena mediterranea</i>	2
Anellida	<i>Amphitrite sp.</i>	2
Cnidaria	<i>Anemonia viridis</i>	2
Anthropoda	<i>Balanus perforatus</i>	2
Mollusca	<i>Bolinus brandaris</i>	2
Tunicata	<i>Bovellus sp.</i>	2
Bryozoa	<i>Bowerbankia sp.</i>	2
Anthropoda	<i>Carcinus maenas</i>	2
Mollusca	<i>Chama gryphoides</i>	2
Mollusca	<i>Chrysalidida sp.</i>	2
Mollusca	<i>Clanculus iussieui</i>	2
Cnidaria	<i>Clyta linearis</i>	2
Mollusca	<i>Columbellula rusticula</i>	2
Cnidaria	<i>Haliplanella luciae</i>	2
Mollusca	<i>Dosinia lupinus</i>	2
Anthropoda	<i>Echinogammarus sp.</i>	2
Mollusca	<i>Ensis siliqua</i>	2
Anellida	<i>Euclymena oerstedii</i>	2
Anellida	<i>Eunice sp.</i>	2
Anellida	<i>Exogaea hebes meridionalis</i>	2
Cnidaria	<i>Garveia franciscana</i>	2
Mollusca	<i>Glaes aculeata</i>	2
Anellida	<i>Glycera tesselata</i>	2
Anellida	<i>Glycera tridactyla</i>	2
Mollusca	<i>Glycimeris insubrica</i>	2
Cnidaria	<i>Gonothryaenae loveni</i>	2
Mollusca	<i>Heleobia stagnorum</i>	2
Mollusca	<i>Hiatella arctica</i>	2
Anthropoda	<i>Hippobole loneirostris</i>	2
Mollusca	<i>Jubibus striatus</i>	2
Anthropoda	<i>Ligia italicica</i>	2
Mollusca	<i>Limaria inflata</i>	2
Mollusca	<i>Macoma cumanina</i>	2
Anthropoda	<i>Mesopodopsis slabberi</i>	2
Mollusca	<i>Musculus marmoratus</i>	2
Mollusca	<i>Natica hebreæ</i>	2
Mollusca	<i>Ocenebrina edwardsi</i>	2
Mollusca	<i>Odostomia sp.</i>	2
Anthropoda	<i>Ophiodromus pallidus</i>	2
Mollusca	<i>Palaeomonetes antennarius</i>	2
Mollusca	<i>Parvicardium minimum</i>	2
Endoprocta	<i>Pedicillina cernua</i>	2
Anthropoda	<i>Periocolades longimanus</i>	2
Cnidaria	<i>Phiatella quadrata</i>	2
Anellida	<i>Prionospio sp.</i>	2
Mollusca	<i>Ptililia sp.</i>	2
Mollusca	<i>Retusa semisulcata</i>	2
Mollusca	<i>Rissoid radiata</i>	2
Cnidaria	<i>Sagartia troyleoides</i>	2
Bryozoa	<i>Scrupocellarria sp.</i>	2
Anellida	<i>Sternaspis scutata</i>	2
Platyhelminthes	<i>Sylochus mediterraneus</i>	2
Anellida	<i>Syllis sp.</i>	2
Mollusca	<i>Tellina donacina</i>	2
Mollusca	<i>Tellina nitida</i>	2
Mollusca	<i>Tellina tenuis</i>	2
Mollusca	<i>Alvania cimex</i>	2
Mollusca	<i>Arca noae</i>	2
Anellida	<i>Arenicola claparedi</i>	2
Anellida	<i>Branchiomma luctuana</i>	2
Bryozoa	<i>Bugula plumosa</i>	2
Anthropoda	<i>Caenis sp.</i>	2
Anellida	<i>Chaetozone setosa</i>	2
Bryozoa	<i>Chorizopora brongniarti</i>	2
Anellida	<i>Cirriformia filigera</i>	2
Tunicata	<i>Clavelina lepadiformis</i>	2
Anellida	<i>Clymenura cylindrica</i>	2
Cnidaria	<i>Cytta haemispherica</i>	2
Anthropoda	<i>Corophium sextonae</i>	2
Anthropoda	<i>Cymodoce truncata</i>	2
Anellida	<i>Dodecaceria concharum</i>	2
Mollusca	<i>Doxax semistriatus</i>	2
Bryozoa	<i>Electra posidonidiae</i>	2
Anthropoda	<i>Pagurus sp.</i>	2
Bryozoa	<i>Fenestrelina malusii</i>	2
Porifera	<i>Halichondria sp.</i>	2
Anthropoda	<i>Iphinoe sp.</i>	2
Anellida	<i>Lentonceris elauca</i>	2
Mollusca	<i>Lithophaga lithophaga</i>	2
Mollusca	<i>Littorina neritoidea</i>	2
Tunicata	<i>Molgula socialis</i>	2
Mollusca	<i>Modiolara subpicta</i>	2
Anthropoda	<i>Pereionotus testudo</i>	2
Mollusca	<i>Petricola lithophaga</i>	2
Anthropoda	<i>Platissa marina</i>	2
Mollusca	<i>Pivinella conica</i>	2
Mollusca	<i>Retusa truncatula</i>	2
Anellida	<i>Scolelepis fuliginosa</i>	2
Bryozoa	<i>Scrupocellarria reptans</i>	2
Porifera	<i>Sycos sp.</i>	2
Mollusca	<i>Montacuta ferruginea</i>	2
Bryozoa	<i>Umbonula ovicellata</i>	2
Anellida	<i>Perinereis macropus</i>	2
Anellida	<i>Syllidia armata</i>	2
Mollusca	<i>Xenostrobus securis</i>	2
Anthropoda	<i>Cleantis prismatic</i>	2
Mollusca	<i>Rissoa sarsi</i>	2
Mollusca	<i>Ventrosia ventrosa</i>	2
Anthropoda	<i>Idotea sp.</i>	2
Bryozoa	<i>Anguineella palmata</i>	2
Bryozoa	<i>Bowerbankia gracilima</i>	2
Cnidaria	<i>Podocoryne carneea</i>	1
Mollusca	<i>Cantharus dorbgini</i>	1
Nematoda	<i>Cyatholamus sp.</i>	1
Anellida	<i>Exosone vergera</i>	1
Bryozoa	<i>Margareta cereoides</i>	1
Anthropoda	<i>Melita sp.</i>	1
Anthropoda	<i>Baetis sp.</i>	1
Anellida	<i>Harmothoe extenuata</i>	1
Anthropoda	<i>Liocarcinus verinalis</i>	1
Porifera	<i>Plakina monolopha</i>	1
Anellida	<i>Siriella clausi</i>	1
Mollusca	<i>Bithynia tentaculata</i>	1
Porifera	<i>Mycale ratalis</i>	1
Mollusca	<i>Brachidontes pharaonis</i>	1
Anthropoda	<i>Chelura turbetra</i>	1
Mollusca	<i>Hypselodoris maculosa</i>	1
Anellida	<i>Nephrys incisa</i>	1
Mollusca	<i>Hydrobia cornica</i>	1
Anthropoda	<i>Nebalia bipes</i>	1
Bryozoa	<i>Bugula simplex</i>	1
Anellida	<i>Nephrys hystericis</i>	1
Anthropoda	<i>Astacilla mediterranea</i>	1
Tunicata	<i>Bovellus leachi</i>	1
Cnidaria	<i>Cordylinophora neapolitana</i>	1
Mollusca	<i>Ensis minor</i>	1
Mollusca	<i>Gibberula philippii</i>	1
Bryozoa	<i>Haplonome graniferum</i>	1
Anellida	<i>Nephrys sp.</i>	1
Anthropoda	<i>Synisona sp.</i>	1
Anellida	<i>Terebellides stroemi</i>	1
Mollusca	<i>Cerithium alucastrum</i>	1
Echinodermata	<i>Genocidaris maculata</i>	1
Anellida	<i>Glyptis propinquus</i>	1
Anellida	<i>Hyalinoecia tubicola</i>	1
Anthropoda	<i>Ischyrocerus inexpectatus</i>	1
Anthropoda	<i>Maera sp.</i>	1
Anellida	<i>Nicolea venustula</i>	1
Anellida	<i>Sabellaria sp.</i>	1
Mollusca	<i>Venus verrucosa</i>	1
Bryozoa	<i>Watersipora complanata</i>	1
Anthropoda	<i>Chironomus sp.</i>	1
Mollusca	<i>Filograna sp.</i>	1
Anthropoda	<i>Gammareopsis sp.</i>	1
Porifera	<i>Geodia cydonium</i>	1
Anellida	<i>Laomone Kroveri</i>	1
Mollusca	<i>Mangelia vaquelinii</i>	1
Porifera	<i>Mycale contarenti</i>	1
Mollusca	<i>Paradoris indecora</i>	1
Anellida	<i>Pileolaria militaris</i>	1
Mollusca	<i>Pusillina sarsi</i>	1
Bryozoa	<i>Scruparia ambigua</i>	1
Anellida	<i>Spirorbanchus polytrema</i>	1
Bryozoa	<i>Victorella muelleri</i>	1
Mollusca	<i>Apnorrhais pespelicani</i>	1
Echinodermata	<i>Astronecten spinulosus</i>	1
Anthropoda	<i>Balanus trigonus</i>	1
Mollusca	<i>Barbatia barbata</i>	1
Cnidaria	<i>Cerianthus membranaceus</i>	1
Mollusca	<i>Cerithium sp.</i>	1
Anellida	<i>Cirratulus sp.</i>	1
Tunicata	<i>Diplosoma listerianum</i>	1
Porifera	<i>Dvidea incrassata</i>	1
Cnidaria	<i>Eudendrium sp.</i>	1
Mollusca	<i>Geodia sp.</i>	1
Anellida	<i>Brania limbata</i>	1
Anellida	<i>Glyptis canepensis</i>	1
Mollusca	<i>Planorbis sp.</i>	1
Anellida	<i>Hydrodides norvegica</i>	1
Mollusca	<i>Hypselodoris elegans</i>	1
Anellida	<i>Leucocarpus chinensis</i>	1
Anthropoda	<i>Limnoria lignorum</i>	1
Anellida	<i>Limnoria sp.</i>	1
Anthropoda	<i>Loxosomella kefersteinii</i>	1
Bryozoa	<i>Muricaria truncata</i>	1
Nematoda	<i>Oncholaimus paralangurenensis</i>	1
Anthropoda	<i>Palaeomon xiphias</i>	1
Anthropoda	<i>Patirambus typicus</i>	1
Phoronida	<i>Phoronis psammophila</i>	1
Anellida	<i>Phyllodocidae</i>	1
Anellida	<i>Pista sp.</i>	1
Anellida	<i>Protula sp.</i>	1
Anellida	<i>Pterocirrus sp.</i>	1
Bryozoa	<i>Schizomavella linearis</i>	1
Bryozoa	<i>Schizomavella linearis v. hastata</i>	1
Mollusca	<i>Scrobicularia rubiginosa</i>	1
Anellida	<i>Tharyx sp.</i>	1
Mollusca	<i>Tricolla speciosa</i>	1
Mollusca	<i>Trinacatella sp.</i>	1
Mollusca	<i>Turbonilla fenestrata</i>	1
Mollusca	<i>Vexillum ebenus</i>	1
Anthropoda	<i>Ampelisca diadema</i>	1
Mollusca	<i>Pisania glabrata</i>	1
Mollusca	<i>Nodulus contortus</i>	1
Mollusca	<i>Anisocyla pointeli</i>	1
Anthropoda	<i>Arcurella dilatata</i>	1
Tunicata	<i>Ascidia conchilega</i>	1
Tunicata	<i>Ascidella aspersa</i>	1
Anthropoda	<i>Ascidia sp.</i>	1
Mollusca	<i>Brachystoma sp.</i>	1
Anellida	<i>Branchiomma sp.</i>	1
Cnidaria	<i>Bunodactis verrucosa</i>	1
Cnidaria	<i>Caltostoma virescens</i>	1
Anellida	<i>Ceratonereis costata</i>	1
Mollusca	<i>Chrysalidida brebreti</i>	1
Anellida	<i>Cirratulus chrysoderma</i>	1
Anellida	<i>Cirrophorus furcatus</i>	1
Cnidaria	<i>Cladocora caespitosa</i>	1
Cnidaria	<i>Cladocoryne floccosa</i>	1
Anthropoda	<i>Clibanarius erythropus</i>	1
Ponera	<i>Cliona sp.</i>	1
Cnidaria	<i>Clivia sp.</i>	1
Mollusca	<i>Corbula sp.</i>	1
Echinodermata	<i>Trachythyre elongata</i>	1
Anellida	<i>Cumella limicola</i>	1
Mollusca	<i>Dendrodoris grandiflora</i>	1
Mollusca	<i>Dentalium inequicostatum</i>	1
Anthropoda	<i>Dexamin spiniventris</i>	1
Cnidaria	<i>Diadumene cincta</i>	1
Mollusca	<i>Didemnum candidum</i>	1
Ponera	<i>Dysidea fragilis</i>	1
Anthropoda	<i>Dyspanopterus savi</i>	1
Bryozoa	<i>Holopoma impressum</i>	1
Bryozoa	<i>Hipnodilus ottomulleriana</i>	1
Anthropoda	<i>Hippolyte leptocerus</i>	1
Mollusca	<i>Hydrobia acuta</i>	1
Mollusca	<i>Hydrobia ulvae</i>	1
Anellida	<i>Hydrodides nigra</i>	1
Ponera	<i>Ircinia variabilis</i>	1
Anellida	<i>Janua sp.</i>	1
Anthropoda	<i>Leander squilla</i>	1
Mollusca	<i>Lepidochiton capreum</i>	1
Platyhelminthes	<i>Leptoplana alcinoi</i>	1
Mollusca	<i>Lineus sp.</i>	1
Anellida	<i>Lumbrineris sp.</i>	1
Mollusca	<i>Lymanea sternalis</i>	1
Anellida	<i>Lysidice ninetta</i>	1
Anthropoda	<i>Melita coroniti</i>	1
Anthropoda	<i>Microdeutopus versicoloratus</i>	1
Mollusca	<i>Mitrella scripta</i>	1
Mollusca	<i>Modiolus adriaticus</i>	1
Mollusca	<i>Mutilus edulis</i>	1
Anthropoda	<i>Nannomya propinqua</i>	1
Mollusca	<i>Nassarius incrassatus</i>	1
Cnidaria	<i>Ocinebrina aciculata</i>	1
Cnidaria	<i>Olinanda phosphorica</i>	1
Nematoda	<i>Oncholaimus campylcercoides</i>	1
Anellida	<i>Opisthodromus avilis</i>	1
Anthropoda	<i>Orchomene humilis</i>	1
Anthropoda	<i>Pasuristes eremita</i>	1
Anellida	<i>Peloscolex sp.</i>	1
Anellida	<i>Pileolaria sp.</i>	1
Mollusca	<i>Pisania striata</i>	1
Mollusca	<i>Platvalvus argo</i>	1
Anthropoda	<i>Potamom fluvatile</i>	1
Bryozoa	<i>Puettina gatvae</i>	1
Anellida	<i>Pulfiella armata</i>	1
Mollusca	<i>Pusia tricolor</i>	1
Mollusca	<i>Puritania purpurea</i>	1

Phylum	Taxon	N. sites	Phylum	Taxon	N. sites	Phylum	Taxon	N. sites
Mollusca	Raphitoma sp.	1	Anellida	Caulieriella caputescoris	1	Arthropoda	Lembos sp.	1
Mollusca	Rissoa albellula	1	Bryozoa	Cellaria fistulosa	1	Anellida	Leptidionotus clava	1
Anellida	Sabellaria alcocci	1	Bryozoa	Cellaria saticornioidea	1	Arthropoda	Leptocheirus guttatus	1
Mollusca	Saccostrea commercialis	1	Bryozoa	Cellepora pumicosa	1	Porifera	Leucetta solida	1
Anellida	Scolelepis tridentata	1	Bryozoa	Cellepora carolinensis	1	Arthropoda	Leucothoe sp.	1
Bryozoa	Scrupocellaria scruposa	1	Bryozoa	Celleporina caminata	1	Bryozoa	Lichenopora radiata	1
Arthropoda	Sirella armata	1	Bryozoa	Celleporina Hassallii	1	Tunicata	Lissoclinum perforatum	1
Anellida	Spirorbis sp.	1	Arthropoda	Ceratopogonidae	1	Arthropoda	Loxoconcha elliptica	1
Cnidaria	Stylactaria sp.	1	Mollusca	Chlamys varia	1		Lumbineris coccinea	1
Anellida	Syllis amica	1	Bryozoa	Chlidonia pyriformis	1	Mollusca	Lunatia sp.	1
Anellida	Syllis cornuta	1	Nematoda	Chromadorella membranata	1	Mollusca	Lymnaea peregra	1
Anellida	Syllis ferruginea	1	Mollusca	Chromodoris sp.	1	Arthropoda	Lystanassa ceratina	1
Bryozoa	Tangamella muelleri	1	Mollusca	Chrysallida manodea	1	Arthropoda	Lystanassa longicornis	1
Arthropoda	Tanysyrum conirostre	1	Arthropoda	Chithamalus montagui	1	Arthropoda	Lystanassa sp.	1
Porifera	Tethya aurantium	1	Arthropoda	Clithamalus stellatus	1	Anellida	Lysidice collaris	1
Nemertina	Tetrasistema sp.	1	Porifera	Clathrina coriacea	1	Mollusca	Mactra glauca	1
Anellida	Thelipus cincinnatus	1	Tunicata	Clavelina phaeogaea	1	Arthropoda	Maera grossimana	1
Mollusca	Tornus subcarinatus	1	Porifera	Cliona viridis	1	Arthropoda	Maera inaequipes	1
Anellida	Tringanovialis rosea	1	Bryozoa	Collarina balzaci	1	Mollusca	Mamilloretusa mamillata	1
Mollusca	Truncatella subcylindrica	1	Bryozoa	Conidozoon planum	1	Anellida	Mediomastus capensis	1
Mollusca	Turritella communis	1	Bryozoa	Puellina radiata	1	Arthropoda	Metaphoxus simplex	1
Endoprocta	Urnatella gracilis	1	Mollusca	Ctena decussata	1	Tunicata	Microcosmus sulcatus	1
Mollusca	Vexillum tricolor	1	Anellida	Ctenodrilus serratus	1	Arthropoda	Microdetonopus obtusatus	1
Porifera	Aaptos aaptos	1	Mollusca	Cylichna crossei	1	Arthropoda	Microdetonopus sp.	1
Mollusca	Acanthochitona fascicularis	1	Arthropoda	Cyprideis torosa	1	Anellida	Microernphytus sphaerocirrata	1
Mollusca	Aclis sp.	1	Nematoda	Cytaholaimus gracilis	1	Bryozoa	Micropora coriacea	1
Mollusca	Acroloxus lacustris	1	Mollusca	Mangelia multifilicola	1	Bryozoa	Micropora ciliata	1
Porifera	Haliclona rosea	1	Nematoda	Daptonema calceolatum	1	Bryozoa	Micropora marsupiata	1
Porifera	Haliclona simulans	1	Platyhelminthes	Dendrocoelum lacteum	1	Arthropoda	Microprotonus maculatus	1
Mollusca	Aequipecten opercularis	1	Mollusca	Dendrodoris limbata	1	Tunicata	Molgeula manhattensis	1
Bryozoa	Aetea sica	1	Tunicata	Didemnum gelatinosum	1	Bryozoa	Mollia patellaria	1
Cnidaria	Aglaoaphenia harpago	1	Tunicata	Didemnum granulosum	1	Mollusca	Monodonta sp.	1
Cnidaria	Aglaoaphenia picardi	1	Mollusca	Dikolens pusilla	1	Mollusca	Monodonida turbinata	1
Cnidaria	Aglaoaphenia plumosa	1	Bryozoa	Disporella hispida	1	Arthropoda	Mumna sp.	1
Cnidaria	Aintasia mutabilis	1	Arthropoda	Dixidae larva	1	Mollusca	Musculus costulatus	1
Cnidaria	Aintasiogiton pellucidus	1	Mollusca	Donacilla cornea	1	Arthropoda	Mysidopsis gibbosa	1
Bryozoa	Alcyonium sp.	1	Mollusca	Donax sp.	1	Arthropoda	Mysis sp.	1
Mollusca	Alvania discors	1	Mollusca	Donax trunculus	1	Arthropoda	Nebalia sp.	1
Mollusca	Alvania pagodula	1	Mollusca	Doriopsis rarispina	1	Anellida	Nematoneurus unicornis	1
Anellida	Amgee adversa	1	Mollusca	Dorvillea rubrovittata	1	Anellida	Nereis kerqueensis	1
Anellida	Amage sp.	1	Anellida	Dorvillea sp.	1	Anellida	Nereis rava	1
Arthropoda	Amnothea hilgendorfi	1	Mollusca	Dosinia sp.	1	Mollusca	Neverita iosephinia	1
Arthropoda	Ampeleis abdita	1	Anellida	Drilonereis filum	1	Mollusca	Nucula nitidosa	1
Arthropoda	Ampeleis sarsi	1	Platyhelminthes	Dugesia sp.	1	Cnidaria	Obelia longissima	1
Arthropoda	Ampeleis sp.	1	Porifera	Dysidea avara	1	Nematoda	Odontophora armata	1
Anellida	Amphictieis gunneri	1	Arthropoda	Echinogammurus stocki	1	Anellida	Odontosyllis duvesiana	1
Arthropoda	Amphithocus brunnneus	1	Arthropoda	Elasmopus pecteniferus	1	Anellida	Odontosyllis gibba	1
Arthropoda	Anthidialia avilis	1	Mollusca	Elysia timida	1	Mollusca	Odostomia plicata	1
Cnidaria	Anthopleura rubripunctata	1	Mollusca	Epitonium commutatum	1	Mollusca	Odostomia scalaris	1
Arthropoda	Aphephusa schiereghenii	1	Anellida	Erpobdella sp.	1	Mollusca	Omalogyra atomus	1
Anellida	Aphrodita aculeata	1	Bryozoa	Escharina vulgaris	1	Bryozoa	Onychocella antiqua	1
Tunicata	Aplidium sp.	1	Anellida	Eudistome pudoritanum	1	Echinodermata	Ophiomyxa pentagona	1
Anellida	Acmira cerruti	1	Anellida	Eulalia macroceros	1	Cnidaria	Orthiphysis asymmetrica	1
Tunicata	Ascidella scabra	1	Anellida	Eulalia punctifera	1	Mollusca	Papillocardium papillosum	1
Mollusca	Ascidolla fragilis	1	Anellida	Eulalia rubinosa	1	Mollusca	Parvicardium vroomi	1
Arthropoda	Asellus aquaticus	1	Anellida	Eulalia trilineata	1	Mollusca	Patella caerulea	1
Echinodermata	Asterina panceri	1	Anellida	Eunice harassi	1	Porifera	Pellina semitubulosa	1
Arthropoda	Athanas nitescens laevirhincus	1	Anellida	Euphosrosea foliosa	1	Mollusca	Peringia elongata	1
Arthropoda	Atyaephyra desmaresti	1	Anellida	Eupolytmia nebulosa	1	Mollusca	Peringia elegans	1
Anellida	Autostolus prolofer	1	Anellida	Exogone naidina	1	Tunicata	Perophora viridis	1
Mollusca	Barlecia unifasciata	1	Bryozoa	Fenestrulina ioannae	1	Anellida	Petaloproticus terricolus	1
Bryozoa	Beania hirtissima	1	Mollusca	Fissurella nubecula	1	Anellida	Pherusa monilifera	1
Bryozoa	Beania magellonica	1	Mollusca	Fusinus syracusanus	1	Bryozoa	Pherussella tubulosa	1
Mollusca	Bela nebula	1	Arthropoda	Gammareopsis maculata	1	Mollusca	Pholads dacyrus	1
Mollusca	Berthella aurantiaca	1	Anellida	Gari costulata	1	Anellida	Anaitides groenlandica	1
Mollusca	Berthella stellata	1	Arthropoda	Gastrosaccus mediterraneus	1	Anellida	Phyllo foetida	1
Mollusca	Bithynia leachii	1	Mollusca	Gibbula umbilicaris	1	Anellida	Pionosyllis sp.	1
Mollusca	Bittinum sp.	1	Mollusca	Gibberula militaris	1	Mollusca	Platycardium papillosum	1
Bryozoa	Bowerbankia imbricata	1	Anellida	Glycera alba	1	Bryozoa	Platyoecia patina	1
Bryozoa	Calanoides venustus	1	Anellida	Glycera rouxi	1	Mollusca	Planorbarius cornutus	1
Anellida	Brania sp.	1	Anellida	Goniada maculata	1	Mollusca	Planorbis planorbis	1
Bryozoa	Caberlea boryi	1	Mollusca	Granularia occulta	1	Anellida	Ptynieretes coccinea	1
Porifera	Cacospongia mollior	1	Mollusca	Gyravulus laevis	1	Cnidaria	Plumularia obliqua	1
Porifera	Cacospongia scalaris	1	Mollusca	Gyravulus sp.	1	Mollusca	Pollia scabra	1
Mollusca	Caecum subannulatum	1	Anellida	Harmothoe spinifera	1	Anellida	Potamilla reniformis	1
Mollusca	Caecum trachea	1	Mollusca	Hemitenton nitidum	1	Arthropoda	Praousellus coxalis	1
Arthropoda	Callinectes danie	1	Echinodermata	Holothuria poli	1	Arthropoda	Processa edulis edulis	1
Arthropoda	Callinectes sapidus	1	Echinodermata	Holothuria (Thymioscygia) impatiens	1	Anellida	Protoarcicia ostreidi	1
Mollusca	Callistostoma kochi	1	Echinodermata	Holothuria sp.	1	Echinodermata	Psammechinus microtuberculatus	1
Bryozoa	Callopista dumerilii	1	Echinodermata	Holothuria tubulosa	1	Anellida	Pseudocapitella incerta	1
Bryozoa	Callopista lineata	1	Mollusca	Hypsodoris fontandrui	1	Arthropoda	Pseudolitris kroverii	1
Mollusca	Calmia glaucoidea	1	Mollusca	Hypsodoris villafranca	1	Mollusca	Pusillina dolium	1
Bryozoa	Calpensia nobilis	1	Arthropoda	Iphimedea minuta	1	Mollusca	Pusillina marginata	1
Mollusca	Calyptraea chinensis	1	Arthropoda	Iphinoe serrata	1	Mollusca	Raphitoma bicolor	1
Anellida	Capitella sp.	1	Porifera	Ircinia spinulosa	1	Bryozoa	Rendantonella violacea	1
Arthropoda	Caprella scaura	1	Arthropoda	Janira maculosa	1	Arthropoda	Rhithropanopeus harrisi	1
Arthropoda	Caprella sp.	1	Cnidaria	Kirchenpauera pinnata	1	Mollusca	Rissoa auriscalpium	1
Arthropoda	Carcinus sp.	1	Cnidaria	Lafeoena tenuis	1	Mollusca	Rissoa guerinii	1
Mollusca	Cassidaria echinophora	1	Mollusca	Lajonkaira lajonkairii	1	Mollusca	Rissoa marginata	1
Anellida	Caulieriella alata	1	Anellida	Laonice cirrata	1	Mollusca	Rissoa membranacea	1

Phylum	Taxon	N. sites	Phylum	Taxon	N. sites	Phylum	Taxon	N. sites
Mollusca	<i>Rissoa paradoxa</i>	1	Bryozoa	<i>Valkeria uva</i>	1	Porifera	<i>Topsisentia genitrix</i>	1
Mollusca	<i>Rissoa pulchella</i>	1	Mollusca	<i>Tapes rhomboides</i>	1	Arthropoda	<i>Heterotanais oerstedi</i>	1
Mollusca	<i>Rissoa similis</i>	1	Mollusca	<i>Venerupis sp.</i>	1	Arthropoda	<i>Echinogammarus mungooides</i>	1
Mollusca	<i>Rissoa opalina</i>	1	Porifera	<i>Anlysina aerophoba</i>	1	Arthropoda	<i>Gammarus crinicornis</i>	1
Nematoda	<i>Sabatieria pulchra</i>	1	Porifera	<i>Verongia sp.</i>	1	Arthropoda	<i>Jassa sp.</i>	1
Angellida	<i>Sabellia sp.</i>	1	Anthropoda	<i>Verrucosa stroemii</i>	1	Arthropoda	<i>Lioxarcinus boliviari</i>	1
Angellida	<i>Sabellides octocirrata</i>	1	Anthropoda	<i>Xestoleberis aurantia</i>	1	Tunicata	<i>Phallusia ingrica</i>	1
Cnidaria	<i>Sarsia sp.</i>	1	Anellida	<i>Orbinia foetida</i>	1	Arthropoda	<i>Iaeropsis sp.</i>	1
Bryozoa	<i>Schizomavella auriculata</i>	1	Mollusca	<i>Diplopontia rotundata</i>	1	Arthropoda	<i>Microdeutopus anomalus</i>	1
Bryozoa	<i>Schizoporella longirostris</i>	1	Anellida	<i>Eumice tortuata</i>	1	Arthropoda	<i>Ampithor sp.</i>	1
Bryozoa	<i>Schizoporella sp.</i>	1	Anthropoda	<i>Jassa falcata</i>	1	Cnidaria	<i>Obelia sp.</i>	1
Anellida	<i>Scalaricia typica</i>	1	Mollusca	<i>Odostomia conoidea</i>	1	Cnidaria	<i>Cordylophora sp.</i>	1
Anellida	<i>Serpula sp.</i>	1	Echinodermata	<i>Ophiactis vires</i>	1	Mollusca	<i>Xenostrobus sp.</i>	1
Bryozoa	<i>Sertella couchii</i>	1	Mollusca	<i>Paludinella littoralis</i>	1	Anellida	<i>Polydora ligni</i>	1
Cnidaria	<i>Sertularella ellisi</i>	1	Anthropoda	<i>Parapenaeus latifrons</i>	1	Anellida	<i>Marpphyza sp.</i>	1
Cnidaria	<i>Sertularia perrusilla</i>	1	Bryozoa	<i>Parasmittina sp.</i>	1	Arthropoda	<i>Sphaeroma sp.</i>	1
Mollusca	<i>Setta semistriata</i>	1	Mollusca	<i>Pinna nobilis</i>	1	Mollusca	<i>Abra sp.</i>	1
Mollusca	<i>Sinex zona cingulata</i>	1	Mollusca	<i>Polyvera dubia</i>	1	Mollusca	<i>Teredo navalis</i>	1
Anthropoda	<i>Sirpus zariquevi</i>	1	Bryozoa	<i>Schizobrachioella sanguinea</i>	1	Tunicata	<i>Didemnum sp.</i>	1
Mollusca	<i>Solen marginatus</i>	1	Bryozoa	<i>Smiliumea reticulata</i>	1	Anthropoda	<i>Gnathia sp.</i>	1
Anellida	<i>Sphaerosyllis hystrix</i>	1	Bryozoa	<i>Watersipora subovoidea</i>	1	Mollusca	<i>Haminoea sp.</i>	1
Anellida	<i>Janua corrugatus</i>	1	Anthropoda	<i>Xantho incisus</i>	1	Mollusca	<i>Bitumia latreillii</i>	1
Porifera	<i>Spongia officinalis</i>	1	Mollusca	<i>Diadora gracea</i>	1	Mollusca	<i>Rissoella globularis</i>	1
Porifera	<i>Spongia virgulosa</i>	1	Anthropoda	<i>Elasmopus affinis</i>	1	Mollusca	<i>Rudolphoseria turriculata</i>	1
Anthropoda	<i>Stenoioea validia</i>	1	Anthropoda	<i>Ammothella longipes</i>	1	Mollusca	<i>Runcina sp.</i>	1
Anthropoda	<i>Stenoioea tergestina</i>	1	Anthropoda	<i>Ischnura elegans</i>	1	Mollusca	<i>Calliostoma laugieri</i>	1
Anellida	<i>Sthenelais poa</i>	1	Anthropoda	<i>Cercion lindeli</i>	1	Mollusca	<i>Mangelia costulata</i>	1
Porifera	<i>Sycon citatum</i>	1	Anthropoda	<i>Hoplodonta sp.</i>	1	Anthropoda	<i>Apseudes sp.</i>	1
Porifera	<i>Sycon raphanus</i>	1	Anthropoda	<i>Cloeon sp.</i>	1	Anthropoda	<i>Paramysis helleri</i>	1
Anellida	<i>Syllides edmona</i>	1	Anthropoda	<i>Gerris sp.</i>	1	Anthropoda	<i>Leucos mediterraneus</i>	1
Anellida	<i>Syllis cirrumpunctata</i>	1	Anthropoda	<i>Notonecta glauca</i>	1	Porifera	<i>Siphonocalima coriacea</i>	1
Mollusca	<i>Tellina fabula</i>	1	Anthropoda	<i>Plea minutissima</i>	1	Mollusca	<i>Setia amabilis</i>	1
Mollusca	<i>Tellina incarnata</i>	1	Anthropoda	<i>Naucoris cimicoides</i>	1	Mollusca	<i>Alvania mammillata</i>	1
Mollusca	<i>Tellina planata</i>	1	Anthropoda	<i>Aeschna mixta</i>	1	Mollusca	<i>Pusillina sp.</i>	1
Mollusca	<i>Tellina pulchella</i>	1	Anthropoda	<i>Lentocerus tineiformis</i>	1	Bryozoa	<i>Beania robusta</i>	1
Nematoda	<i>Terschellingia communis</i>	1	Anthropoda	<i>Agraylea sexmaculata</i>	1	Anellida	<i>Pseudoleiocapitella fauveti</i>	1
Nematoda	<i>Terschellingia longicaudata</i>	1	Anthropoda	<i>Argyroreta aquatica</i>	1	Anellida	<i>Cirratulus filiformis</i>	1
Mollusca	<i>Thais haemastoma</i>	1	Anthropoda	<i>Corixa affinis</i>	1	Anellida	<i>Brania pusilla</i>	1
Mollusca	<i>Thracia corbuloides</i>	1	Anthropoda	<i>Sympetrum sp.</i>	1	Anellida	<i>Marpphyza bellii</i>	1
Mollusca	<i>Thysas flexuosa</i>	1	Anthropoda	<i>Chaoborus plumicornis</i>	1	Mollusca	<i>Bithynella sp.</i>	1
Anthropoda	<i>Titanethes albifus</i>	1	Bryozoa	<i>Cheilostomata onusca</i>	1	Mollusca	<i>Retusa sp.</i>	1
Mollusca	<i>Turbonilla delicata</i>	1	Bryozoa	<i>Cheilostomata osconhora</i>	1	Plathelminthes	<i>Sylvocus pilidium</i>	1
Mollusca	<i>Turbonilla lactea</i>	1	Bryozoa	<i>Tubulipora flabellaris</i>	1	Anthropoda	<i>Eriphia spinifrons</i>	1
Mollusca	<i>Turbonilla obliquata</i>	1	Mollusca	<i>Gibbula richardi</i>	1	Anthropoda	<i>Callianassa candida</i>	1
Mollusca	<i>Turbonilla striatula</i>	1	Echinodermata	<i>Holoturia polii</i>	1	Anthropoda	<i>Sacculina sp.</i>	1
Mollusca	<i>Turritella turbona</i>	1	Anellida	<i>Streblospio dekhuzeni</i>	1			